


ENDOCRINE AND IMMUNE PROFILES OF IMMATURE PINNIPEDS


By

Mandy Jean Keogh


RECOMMENDED:



Dr. Michael Castellini


Dr. Tanya Holliman


Dr. Rudy Ortiz


Dr. Jonathan Runstadler



Dr. Shannon Atkinson, Advisory Committee Chair



Dr. Katrin Iken
Head, Program in Marine Science and Limnology

APPROVED:


Dr. Michael Castellini

Dean, School of Fisheries and Ocean Sciences


Dr. Lawrence Duffy, Dean of the Graduate School


Date July 11, 2011

ENDOCRINE AND IMMUNE PROFILES OF IMMATURE PINNIPEDS

A

DISSERTATION

Presented to the Faculty
of the University of Alaska Fairbanks

in Partial Fulfillment of the Requirements
for the Degree of

DOCTOR OF PHILOSOPHY

By

Mandy Jean Keogh, B.S., M.S.

Fairbanks, Alaska

August 2011

UMI Number: 3484671

All rights reserved

INFORMATION TO ALL USERS

The quality of this reproduction is dependent upon the quality of the copy submitted.


In the unlikely event that the author did not send a complete manuscript and there are missing pages, these will be noted. Also, if material had to be removed, a note will indicate the deletion.



UMI 3484671

Copyright 2011 by ProQuest LLC.

All rights reserved. This edition of the work is protected against unauthorized copying under Title 17, United States Code.



ProQuest LLC
789 East Eisenhower Parkway
P.O. Box 1346
Ann Arbor, MI 48106-1346

Abstract

There is increasing interest in assessing the health of individuals and populations of pinnipeds found in the North Pacific, primarily due to population declines leading to conservation concerns. This study assessed the “health” of animals by quantifying hormones associated with fat mass (leptin), lipid and water metabolism (cortisol and aldosterone), and growth and metabolism (thyroxine and triiodothyronine) as well as circulating total and differential leukocyte counts and *in vitro* proliferation of peripheral blood mononuclear cells (PBMC). Body mass and condition are influenced by an individual’s disease and nutritional state. Glucocorticoids are known to affect the immune system and may be stimulated by a multitude of factors. I hypothesized that age or body mass would influence leukocyte counts, PBMC proliferation, and hormone concentrations in Steller sea lion (*Eumetopias jubatus*) pups and that the response of cortisol to an acute stressor would impact immune parameters in juvenile harbor seals (*Phoca vitulina*). Further, given the inherent requirements of disturbance and animal handling necessary for sampling pinnipeds, the impact of these activities on endocrine and immune profiles was assessed.

Total white blood cell (WBC) counts, neutrophil counts and T cell proliferation decreased with increasing age in Steller sea lion pups. However, no relationship between body condition index and circulating concentration of hormones quantified was detected. Circulating concentrations of cortisol, thyroxine, and triiodothyronine were influenced by the rookery disturbance. However, the variation attributed to the disturbance was low and did not alter total or differential WBC counts or *in vitro* proliferation of PBMC. In harbor seals, cortisol and aldosterone concentrations increased following an acute stressor which resulted in a stress leukogram. Total WBC decreased driven primarily by the decrease in neutrophil counts with simultaneous increase in lymphocytes leading to an overall decrease in neutrophil to lymphocyte ratio. These findings highlight the endocrine system’s influence on the immune system in immature pinnipeds.

Table of Contents

	Page
Signature Page	i
Title Page	ii
Abstract	iii
Table of Contents	iv
List of Figures	viii
List of Tables	x
List of Appendices	xi
List of Abbreviations	xii
Acknowledgements	xiv
CHAPTER 1: INTRODUCTION	1
1.1 Overview	1
1.1.1 Objectives	3
1.2 Life History of Pinnipeds	4
1.2.1 Steller Sea Lions	5
1.2.2 Harbor Seals	7
1.3 Body Condition	8
1.4 Endocrine System	11
1.4.1 Hypothalamic-Pituitary-Adrenal Axis	11
1.4.2 Thyroid Gland and Hormones	13
1.4.3 Leptin	14
1.5 Immune System	15
1.5.1 Immune Organs and Leukocytes	15
1.5.2 Hematology and PBMC Proliferation	18
1.6 Literature Cited	20

CHAPTER 2: Steller sea lion (<i>Eumetopias jubatus</i>) pups undergo a decrease in circulating white blood cells and the ability of T cells to proliferate during early post-natal development	38
2.1 Abstract	38
2.2 Introduction.....	38
2.3 Materials and Methods.....	40
2.3.1 Study Site and Animals.....	40
2.3.2 Complete Blood Cell Count.....	42
2.3.3 Cortisol Radioimmunoassay	42
2.3.4 PBMC Isolation and Proliferation	42
2.3.5 Statistical Analysis.....	44
2.4 Results and Discussion	44
2.5 Acknowledgements.....	50
2.6 Literature Cited	50
CHAPTER 3: Body condition and endocrine profiles of healthy Steller sea lion (<i>Eumetopias jubatus</i>) pups during the early postnatal period.....	56
3.1 Abstract	56
3.2 Introduction.....	57
3.3 Materials and Methods.....	59
3.3.1 Animals and Body Condition.....	59
3.3.2 Blood Collection	59
3.3.3 Hormone Radioimmunoassays	60
3.3.4 Statistical Analysis.....	60
3.4 Results.....	61
3.4.1 Age and Body Mass	61
3.4.2 Hormone Profiles and Relationships	61
3.5 Discussion	68
3.6 Acknowledgements.....	72

	Page
3.7 Literature Cited	73
CHAPTER 4: Endocrine and immunological responses to adrenocorticotrophic hormone (ACTH) infusion in juvenile harbor seals (<i>Phoca vitulina</i>) during winter and spring.....	78
4.1 Abstract	78
4.2 Introduction	79
4.3 Material and Methods	81
4.3.1 Animals and Blood Collection	81
4.3.2 Hormone Radioimmunoassays	82
4.3.3 Total and Differential Leukocyte Counts	83
4.3.4. PBMC Isolation and Proliferation	83
4.3.5 Statistical Analysis	85
4.4 Results	85
4.4.1 Changes in Hormone Concentrations Following ACTH	85
4.4.2 Alterations in Total and Differential Leukocyte Counts	89
4.4.3 PBMC <i>in vitro</i> Proliferation	89
4.5 Discussion	92
4.5.1 Hormone Concentrations Induced by ACTH	92
4.5.2 Alterations in Total and Differential Leukocyte Counts	94
4.5.3 PBMC <i>in vitro</i> Proliferation	95
4.5.4 Body Mass	97
4.6 Conclusions	97
4.7 Acknowledgements	98
4.8 Literature Cited	98
CHAPTER 5: General Conclusions	106
5.1.1 Hormones, Body Mass and Age	106
5.1.2 Total and Differential Leukocyte Counts	108
5.1.3 PBMC Proliferation	110
5.1.4 Effect of Handling and Sampling	111

	Page
5.2 Literature Cited	111
Appendices.....	117

List of Figures

	Page
Figure 2.1. PBMC proliferation responses from 21 SSL pups exposed to 1.0 µg/ml ConA. Results presented as (A) absorbency (OD) and (B) stimulation index (SI). Significant ($p \leq 0.05$) trend lines are presented	46
Figure 2.2. (A) Total WBC counts (103/µl) from 42 pups during the 38 days following birth and differential counts for (B) neutrophils (103/µl); (C) lymphocytes (103/µl); (D) monocytes (103/µl); and (E) eosinophils (103/µl). (F) Total WBC counts (103/µl) (G) neutrophils (103/µl); (H) lymphocytes (103/µl); (I) monocytes (103/µl); and (J) eosinophils (103/µl) are also presented against level of circulating cortisol (ng/ml). Significant ($p \leq 0.05$) trend lines are presented....	49
Figure 3.1. A total of 61 pups (male=37, female=24) were sampled during 2005, 2007, and 2008. Morphometrics for individual pups at age are presented in (a) body condition index, (b) axillary girth, (c) standard length, and (d) body mass. Female pups are represented by (●) and male pups are by (○).	62
Figure 3.2. (a) Aldosterone concentrations (pg/ml) and (b) cortisol concentrations (ng/ml) from female (●) and male (○) SSL pups. Dashed line separates the pups with greater than 2 SD of the mean for male and female pups for aldosterone concentration (pg/ml).....	64
Figure 3.3. Circulating concentrations of (a) cortisol (ng/ml); (b) total T4 (ng/ml); (c) free T4 (pg/ml); (d) total T3 (ng/ml) from female (●) and male (○) SSL pups regressed against the elapsed time from the initial arrival on the rookery to the time when individual pups were removed from the corral for blood collection. Female pups are represented by filled circles, males are by hollow circles. Significant ($p \leq 0.05$) trend lines are presented	66

- Figure 3.4. Circulating thyroid hormone concentrations from female (●) and male (○) SSL pups. (a, b) total T4 (ng/ml); (c, d) free T4 (pg/ml); (e, f) total T3 (ng/ml) presented against the body condition index (BCI) or age. Female pups are represented by filled circles, males are by hollow circles. Only significant ($p \leq 0.05$) trend lines are presented.67
- Figure 4.1. Mean (\pm SD) serum cortisol (A) and aldosterone (C) concentrations over time following ACTH injection during the winter (●) and spring (○) for seven juvenile harbor seals ($n=7$). Area under the curve (AUCI) following ACTH injection for cortisol (B) and aldosterone (D) during the winter and spring experiments. * denotes significant ($p < 0.05$) difference from time 0.88
- Figure 4.2. Mean (\pm SD) serum concentration for total (A) and free (B) thyroxine and total triiodothyronine (C) following ACTH injection during the winter (●) and spring (○) for seven juvenile harbor seals ($n=7$). Dashed lines denote baseline concentrations.90
- Figure 4.3. Mean differential leukocyte counts for all seven juvenile harbor seals following an acute stressor (exogenous ACTH) during the winter with results reported as percentage (A) and absolute count (B) and during spring with results reported as percentage (C) and absolute count (D).....91
- Figure 4.4. PBMC proliferation responses from seven captive juvenile harbor seals exposed to (A) 1.0 $\mu\text{g/ml}$ ConA and (B) 100 $\mu\text{g/ml}$ LPS following an acute stressor during winter (●) and spring (○). Results presented as mean stimulation index (SI) \pm SD92

List of Tables

	Page
Table 2.1. Mean (\pm SD) for age, body mass (BM), standard length (SL), and axillary girth (AG) for SSL pups by sex and year.	41
Table 2.2. Mean (\pm SD) and range of cortisol, hematology and PBMC proliferation from SSL pups. Age effect was significant at $p < 0.050$	45
Table 3.1. Mean (\pm SD) for age, body mass, standard length, and axillary girth for SSL pups by year and sex.	63
Table 3.2. Mean (\pm SD) for cortisol, aldosterone, total and free thyroxine (T4), total triiodothyronine (T3), and leptin concentrations for all 61 SSL pups and separated by sex (female=24, male=37).	65
Table 4.1. Basal (time 0) mean (\pm SD) and range of cortisol, aldosterone, total and differential blood cell counts and PBMC proliferation by season from captive juvenile harbor seals (n=7).	87

List of Appendices

	Page
Appendix A. List of Commercially Available RIA and Reagents.....	117
Appendix B. Standard Operating Protocol for Aldosterone Assays.....	118
Appendix C. Standard Operating Protocol for Cortisol Assays	119
Appendix D. Standard Operating Protocol for Total Thyroxine (T ₄) Assays.....	120
Appendix E. Standard Operating Protocol for Free Thyroxine (T ₄) Assays	121
Appendix F. Standard Operating Protocol for Total Triiodothyronine (T ₃) Assays.....	122
Appendix G. Standard Operating Protocol for Leptin Assays.....	123
Appendix H. Standard Operating Protocol for Cell Preparation Tubes (CPT™).....	125
Appendix I. Standard Operating Protocol for Sodium Heparin Tubes	126
Appendix J. Standard Operating Protocol for PBMC Proliferation Assays	127
Appendix K. Solutions.....	129

List of Abbreviations

ACTH	adrenocorticotrophic hormone
AG	axillary girth
ASLC	Alaska SeaLife Center
BCI	body condition index
BM	body mass
BrdU	bromodioxuridine
CBC	complete blood cell
CASE	Changing Alaska Science Education
ConA	concanavalin A
CPT	cell preparation tube
DMSO	dimethyl sulfoxide
DPS	distinct population segments
ETOH	ethanol
FBS	fetal bovine serum
GC	glucocorticoid
Hb	hemoglobin
Hct	hematocrit
HPA	hypothalamic-pituitary-adrenal axis
HS	harbor seal
HE	human equivalent
ISO	isoflurane
LPS	lipopolysaccharide
MCHC	mean corpuscular hemoglobin concentration
MCV	mean corpuscular volume
NFS	northern fur seals
NZSL	New Zealand sea lions

OD	optical density
OC	organochlorine
PBMC	peripheral blood mononuclear cells
PBS	phosphate buffered saline
PCCRC	Pollock Conservation Cooperative Research Center
Plt	platelet counts
PR	physically restrained
RIA	radioimmunoassay
RBC	red blood cell
SL	standard length
SSL	Steller sea lion
SI	stimulation index
T ₄	thyroxine
T ₃	triiodothyronine
WBC	white blood cell

ACKNOWLEDGEMENTS

I would like to thank my committee members for their encouragement and help with this project. I wish to thank Dr. Shannon Atkinson for always being open to my ideas and for the opportunity to work in her lab and explore Alaska, both of which I have tried to take full advantage of. Dr. Michael Castellini's excitement for research and fresh perspective of the data was greatly appreciated and helped me with the big picture of this dissertation and my career; Dr. Tuula Hollmen provided guidance and suggestions for data analysis and only booed me once for sounding the door alarm near her office; I greatly appreciate Dr. Rudy Ortiz for his edits and critical reviews and most importantly for constantly pushing and challenging me as a researcher; Dr. Jonathan Runstadler was incredibly supportive offering seminars by distance in immunology and infectious disease, expanding my exposure to topics beyond the narrow scope of this dissertation.

Funding for this project was provided by the Alaska SeaLife Center (ASLC) with congressionally appropriated funds provided by National Marine Fisheries Service and a grant awarded to S. Atkinson by the Pollock Conservation Cooperative Research Center (PCCRC). However, the findings and conclusions presented by the author are my own and do not necessarily reflect the views or positions of the Center or the University of Alaska. Additional funding was provided through a fellowship from the National Science Foundation GK-12 award to the University of Alaska Fairbanks "New GK-12 Program: The CASE (Changing Alaska Science Education) for Enhancing Understanding of Climate Change". Travel funds were provided by the ASLC, PCCRC, the Matt Myers memorial graduate student travel award, and student travel awards from the International Association of Aquatic Animal Medicine and the Society for Marine Mammalogy, enabling me to attend and present this research at several national and international conferences.

The staff at the Seward Marine Center, the School of Fisheries and Ocean Sciences and the ASLC provided support enabling the completion of the majority of this

project while at a remote campus. Conducting this project at the ASLC afforded me several opportunities to participate in unique research and field settings. I am especially grateful to John Maniscalco for the opportunity to collaborate with the Chiswell Island remote monitoring program and to Dr. Jo-Ann Mellish for letting me tag along on capture cruises and work up days for the Transient Juvenile Steller sea lion project. Dr. Russ Andrews attempted to provide samples while Dr. Lori Polasek facilitated samples from the captive harbor seals and numerous discussions, both of which were appreciated. This project benefitted from the support and expertise of the husbandry and veterinary staff at the ASLC including Jane Belovarac, Jen Dailer, Heather Down, Millie Gray, Lisa Hartman, Lee Keller, Aaron Lacy, Lynda Leppert, Brett Long, Elizabeth Moundalexis, Megan Peterson, Christy Phillips, Amber Stephens, Carol Stephens, and Drs. Pam Tuomi and Carrie Goertz.

At the onset of my project I was fortunate to have numerous graduate students located in Seward to help with studying at a remote campus, making courses by distance less intimidating. A special thank you to my fellow graduate students and lab mates; Beate Litz, Abbie Elsworth, Brook McFarland, Jared Guthridge, Jason Waite, Matt Myers, Jeanette Nienaber, Rebekka Federer, Sara Carroll, Mary Bozza, Lisa Petrauskas, and Jacqueline Ramsay. I have fully enjoyed my time in Alaska and have made incredible friends who helped me in my attempt to balance graduate school and the rest of my life. I am grateful for the numerous hiking, skiing and weekend trips exploring Alaska with Ann Riddle, Jill Prewitt, Lance and Cheryl Clark, Shiway Wang, Dave Safine, John Skinner, Amber Stephens, Rebekka Federer, Jeanette Nienaber, Abbie Ellsworth, Pam Parker, and the many dogs that accompanied us.

I am incredibly fortunate to have a large and boisterous family all of whom have been more than just supportive, but rather outright excited about whatever I was doing. My parents have always provided me with support, encouragement and many frequent flier miles, which have been instrumental in my long and seemingly unending academic career. I am grateful for my brothers Eric, Keith and Jason for simply being there for as long as I can remember. My sister-in-law Anne and my nephews Griffin, Carson, and

Weston have been great additions to the family. My pets, Libby, Emma, and Cali were constant companions and sources of entertainment and unfortunately some large veterinary bills (Emma and Cali) and several car repairs (Libby). I am indebted to numerous people for watching my pets while I traveled including Amber Stephens, Ann Riddle, Jill Prewitt, Jason Waite, Brenna McConnell, David Booher and especially Jeanette Nienaber.

Chapter 1: Introduction

1.1 Overview

There is increasing interest in assessing the health of individuals and populations of North Pacific pinnipeds primarily due to recent population declines leading to conservation concerns (Castellini *et al.*, 1993; Zenteno-Savin *et al.*, 1997; Rea *et al.*, 1998; Trumble and Castellini, 2001; Trumble and Castellini, 2002; Burek *et al.*, 2003; Burek *et al.*, 2005; Goldstein *et al.*, 2007; Atkinson *et al.*, 2008). The decline in pinniped populations has coincided with changing environmental conditions of natural and anthropogenic origin (Le Boeuf and Crocker, 2005; Guenette *et al.*, 2006; Atkinson *et al.*, 2008; Burek *et al.*, 2008). Several species of marine mammals and birds in the Northeast Pacific have been in decline since the mid-1970's (Pitcher, 1990; Loughlin, 1998; Small *et al.*, 2003; Jemison *et al.*, 2006). The number and diversity of species that experienced a decline in the Northeast Pacific led to various hypotheses based on shared ecological factors across taxa such as predation by killer whales (*Orcinus orca*), shifts in ocean productivity as a result of the Pacific Decadal Oscillation, competition with fisheries, and interspecific competition for resources.

Affected pinniped species and the focus of this dissertation include Steller sea lions (SSL; *Eumetopias jubatus*) and harbor seals (HS; *Phoca vitulina*). Population trends of HS have varied with most stocks currently being stable or increasing in numbers (Harvey *et al.*, 1990; Jemison *et al.*, 2006). However, within Alaska several populations are still reduced compared to numbers in the 1970 and 1980's (Pitcher, 1990; Frost *et al.*, 1999; Hoover-Miller *et al.*, *in press*), and specifically within Prince William Sound the harbor seal population has continued to decrease (Ver Hoef and Frost, 2003). Currently, two distinct population segments (DPS) of SSL are recognized within U.S. waters: the eastern and western DPS. The western DPS is listed as endangered under the Endangered Species Act (U.S. Federal Register 62:30772–30773), having declined by more than 80% since the 1970's (Loughlin, 1998; Sease *et al.*, 2001). The cause of the population decline

and/or failure to recover of HS and SSL has been the focus of continuing research and debate (Springer *et al.*, 2003; DeMaster *et al.*, 2006; Trites *et al.*, 2007a; Atkinson *et al.*, 2008). Guenette *et al.* (2006) concluded that at different spatial and temporal scales four factors including predation by killer whales, shifts in ocean productivity as a result of the Pacific Decadal Oscillation, competition with fisheries, and interspecific competition for resources likely played a significant role in the population trends of SSL. These factors would also likely influence the population trends of other species in the same ecosystem. Excluding predation by killer whales, these factors have been proposed to lead to reduced body condition, mass or size of individuals potentially affecting the health of individuals and populations. However, there remains no clear reason for the observed decline.

Changes in body condition or mass have been associated with changes in circulating hormones in several species of pinnipeds (Jeanniard du Dot *et al.*, 2009; Ortiz *et al.*, 2003b). The neuroendocrine system influences immune responses through neuronal and hormonal signals originating from the autonomic nervous system, hypothalamic-pituitary-adrenal axis (HPA), and the renin-angiotensin-aldosterone system. Conversely, the immune system feeds back to the sympathetic/parasympathetic nervous system and the HPA axis. These finely tuned interactions are required for the survival and health of an individual and are influenced by body condition, reproductive status, and exposure to stressors (Haddad *et al.*, 2002; Weber, 2003). The mammalian immune system protects against foreign pathogens and plays a vital role in the repair of injured tissues and therefore perturbations may lead to disease. Further, seasonally breeding mammals commonly exhibit significant annual cycles in body mass and energy demands (Lincoln and Richardson, 1998; Rousseau *et al.*, 2003). These changes in body mass and/or nutritional status are characterized by alterations in concentrations of several hormones that influence body condition (Bhat *et al.*, 2003). Similar seasonality occurs in pinnipeds (Rosen and Renouf, 1997; Rosen and Renouf, 1998; Kumagai *et al.*, 2006; Mellish *et al.*, 2007; Rosen and Kumagai, 2008), which has been associated with changes in cortisol and thyroid hormones (Riviere *et al.*, 1977; Ashwell-Erickson *et al.*, 1986;

Boily, 1996; Agarwal and Marshall, 1998; Oki and Atkinson, 2004; Rosen and Kumagai, 2008).

This study assesses the “health” of individual animals by quantifying circulating concentrations of hormones associated with nutritional status and fat mass (leptin), lipid and water metabolism (cortisol and aldosterone), and growth and metabolism (thyroxine and triiodothyronine) as well as circulating total and differential leukocyte counts, and by assessing the *in vitro* proliferation of peripheral blood mononuclear cells (PBMC). Body mass and composition are influenced by an individual’s disease and nutritional state and may act as a proxy for assessing an individual’s or population’s health. Therefore, age and body mass (or condition) is proposed to be an important factor in the health of juvenile HS and SSL pups. I hypothesize that age or body size will be associated with: 1) circulating leukocyte counts; 2) peripheral blood mononuclear cell proliferation; 3) baseline hormone concentrations; and 4) changes in hormone concentrations and immune parameters in response to exogenous adrenocorticotrophic hormone (ACTH) stimulation.

1.1.1 Objectives

Objective 1: Evaluate the influence of age on the total and differential leukocyte counts and *in vitro* proliferation of PBMC in SSL pups during the early post-natal period.

Objective 2: Evaluate the influence of body condition on circulating concentrations of hormones associated with nutritional status and fat mass (leptin), lipid and water metabolism (cortisol and aldosterone), and growth and metabolism (thyroxine and triiodothyronine) during the early postnatal period in SSL pups.

Objective 3: Determine the impact of rookery disturbance and animal handling necessary for sampling wild pinnipeds on endocrine (cortisol, aldosterone, thyroxine, triiodothyronine, and leptin) and immune (leukocyte counts and *in vitro* proliferation of PBMC) profiles of SSL pups.

Objective 4: Evaluate the physiological response to an acute stressor (exogenous ACTH) on endocrine (cortisol, aldosterone, thyroxine, and triiodothyronine

concentrations) and immune parameters (total and differential leukocyte count and *in vitro* proliferation of PBMC) in juvenile HS.

1.2 Life History of Pinnipeds

Pinnipeds are a monophyletic group having diverged from a shared arctoid carnivore ancestor that reentered the sea approximately 25 million years ago (Berta and Adam, 2001) and subsequently diverged into three families: Otariidae (sea lions and fur seals), Odobenidae (one extant species, the walrus), and Phocidae (true seals). All extant pinnipeds are amphibious, dividing their time between foraging at sea and hauling out on land or sea ice to reproduce, molt, and rest. There is striking diversity in life history traits (e.g. mating and lactation strategies, body size, degree of sexual dimorphism) between and within the three pinniped families, likely driven by multiple factors including thermoregulation, distribution of resources, predation pressures, breeding substrate and phylogenetic constraints (Costa, 1991; Boness and Bowen, 1996; Boyd, 1998).

In general pinnipeds are aggregate, seasonal breeders with mating occurring in water or on land (or ice) resulting in highly synchronized terrestrial parturition of a single pup. Gestational periods are relatively long ranging between 9.5 and 16 months of which 1.5-5 months is an embryonic diapause (Bowen, 1991). Adult female phocids are generally larger than otariids resulting in phocid pups having larger absolute mass at birth and faster postnatal growth rates compared to otariid pups (Bowen, 1991; Costa, 1991; Kovacs and Lavigne, 1992). When compared on a mass-specific basis of the adult female, mass of pups at birth are similar between phocids and otariids (Kovacs and Lavigne, 1986; Kovacs and Lavigne, 1992) while otariid pups are larger (and older) at weaning than phocids (Bowen, 1991; Costa, 1991). As with all placental mammals, supporting the nutritional needs of an offspring through weaning is the period of the highest energetic cost of reproduction (Millar, 1977).

Though overly simplistic, several generalizations are often made about the life history strategies of phocids and otariids. Shorter lactation periods ranging between 4-50 days (Bowen, 1991) are found in phocids and are characterized by fat content in the milk

between 40-60% (Schulz and Bowen, 2004) and a larger energy store in the form of a subcutaneous blubber layer in the adult female phocid (per kilogram of body mass) compared to otariid (Costa, 1991). This large blubber layer supports both milk production and the nutritional requirements of the adult female during the fasting period associated with nursing. The pups are abruptly weaned when the adult female leaves the pup in order to forage, though some smaller phocids such as the harbor seal forage while supporting the nutritional needs of their pups (Boness *et al.*, 1994). The large blubber layer of the pup, which is acquired during the relatively short nursing period, provides the nutritional support for the pup during its post weaning fast and while it develops swimming and foraging skills (Bowen, 1991; Mellish *et al.*, 1999; Frost *et al.*, 2006; Clark *et al.*, 2007).

The relatively longer lactation period for otariids of 17-77 weeks are characterized by fat content in milk ranging between 14-44% (Arnould *et al.*, 1996; Boness and Bowen, 1996; Arnould and Hindell, 1999) and smaller subcutaneous blubber layer in the adult female. Otariids have “foraging cycle strategy” of alternating forage bouts between nursing periods in order to meet their own nutritional needs as well as those of the pup. Otariid pups remain at the rookery, fasting until the adult female returns from foraging trips (Bonner, 1984; Oftedal *et al.*, 1987; Trillmich, 1990; Costa, 1991). Foraging trips will last hours to days and increase in duration as the pup matures (Maniscalco *et al.*, 2006). The pup will begin to forage with the adult female to supplement its nutritional needs and is gradually weaned, generally prior to the birth of the new pup.

The current study focuses on two species: one otariid, the SSL and one phocid, the Pacific HS. Therefore, the remainder of this section will focus on the specific life history strategies and conservation status of these two species.

1.2.1 Steller Sea Lions

The SSL is the largest otariid and exhibits dramatic sexual dimorphism (male’s annual average body mass 556 kg, breeding season 1,120 kg; females’ annual average

body mass 263 kg, breeding season 350 kg). Similar to other otariids, SSL have seasonal and highly synchronized parturition and mating, which requires a return to land. This reproductive strategy of terrestrial parturition is characterized by large aggregations of territorial bulls, adult females, yearlings and pups present seasonally on the rookery.

At the time of birth, pups remain with the adult female during the perinatal period, ranging between 1.7 to 17.1 days (Merrick *et al.*, 1987; Hood and Ono, 1997; Maniscalco *et al.*, 2006). Pups have continuous access to the female for nursing during the perinatal period, which ends when the female returns to the sea for the first foraging trip since giving birth, leaving the pup unattended on land for several hours to days. For the remainder of the lactation period the female will alternate between short foraging trips to sea and visits to land to nurse the pup. The lactation period lasts about one year during which the female may be pregnant with next year's pup.

SSL are distributed along the North Pacific Rim from northern Japan to California (Loughlin *et al.*, 1984). Within U.S. waters there are currently 2 federally recognized distinct population segments (DPS): the eastern and western DPS, divided along the 144° W longitude (Cape Suckling, Alaska), based upon variation in mitochondrial control-regions (Bickham *et al.*, 1996; Bickham *et al.*, 1998). The eastern stock includes rookeries in California, Oregon, British Columbia, and southeastern Alaska while the western stock is comprised of rookeries within Prince William Sound, the Bering Sea, Central and Western Gulf of Alaska, the Aleutian Islands, Russia, and the Kuril Islands. Within U.S. waters, both DPS fall under the jurisdiction of the NOAA Fisheries with the western DPS currently listed as endangered and the eastern DPS as threatened under the Endangered Species Act and depleted throughout its range under the Marine Mammal Protection Act.

The SSL population has declined by more than 80% since the 1970's (Sease *et al.*, 2001). There are several potential factors that may have led to the decline and/or lack of recovery including predation, disease, contaminants, and nutritional stress (Atkinson *et al.*, 2008). While the mechanistic cause of the decline is a matter of continuing debate (Springer *et al.*, 2003; Fritz and Hinckley, 2005; DeMaster *et al.*, 2006; Trites *et al.*,

2007b; Atkinson *et al.*, 2008), several studies have suggested a reduction in juvenile survival (York, 1994; Holmes and York, 2003) and a long-term decrease in natality rates (Holmes and York, 2003; Holmes *et al.*, 2007). In spite of the dramatic population decline, Baker *et al.*, (2005) found the diversity of haplotypes and nucleotides within the mitochondrial control region were high and moderate, respectively, across the entire range of SSL, concluding that there was no evidence of a genetic bottleneck.

1.2.2 Harbor Seals

HS are currently classified into four recognized subspecies found in either the Atlantic (North America: *Phoca vitulina concolor*; Europe: *Phoca vitulina vitulina*) or Pacific Oceans (North America: *Phoca vitulina richardsii*; Asia: *Phoca vitulina stejnegeri*). A fifth subspecies, based upon morphology and behavior has been proposed for a freshwater population found in Quebec, Canada (*Phoca vitulina mellonae*) (Smith and Lavigne, 1994). The classification of two subspecies in the Pacific HS was originally based on morphological differences including body size, coat color and cranial morphology (Temte *et al.*, 1991). However, more recent molecular studies have not supported the division of the Pacific HS into two subspecies (Westlake and O'Corry-Crowe, 2002).

Phoca vitulina concolor and *Phoca vitulina richardsii* can be found coastally within U.S. waters of the Atlantic and Pacific Oceans, respectively. For management purposes NMFS currently divides the two subspecies into seven stocks including one Atlantic (Western North Atlantic) and six Pacific (Bering Sea, California, Gulf of Alaska, Oregon-Washington coastal, Southeast Alaska, and Washington Inland). Population trends have varied for the different management stocks with most stocks stable or increasing since the Marine Mammal Protection Act was enacted in 1972 (Harvey *et al.*, 1990; Jemison *et al.*, 2006). However, the Gulf of Alaska stock may be continuing to decline and is still reduced compared to population numbers in the 1970 and 1980's (Pitcher, 1990; Frost *et al.*, 1999; Hoover-Miller *et al.*, *in press*) and specifically in Prince William Sound (Ver Hoef and Frost, 2003) while other populations in southeastern

Alaska, excluding Glacier Bay which has declined (Mathews and Pendleton, 2006), have increased (Small *et al.*, 2003).

HS are small phocids displaying some sexual dimorphism with adult male (average body mass 115 kg; average length 175 cm) being larger than female harbor seals (80 kg; 160 cm). They are polygamous, aquatic breeders with males showing plasticity in mating behaviors, which often involves occupying underwater territories where they perform vocal displays for females (Bowen *et al.*, 2006). Females have been observed to first give birth between 4 and 6 years of age and continue to grow after sexual maturity until about 10 years (Markussen *et al.*, 1989). This has led to numerous studies to assess the effect of maternal body condition and age on birth and weaning mass and the survival of the pup (Ellis *et al.*, 2000; Bowen *et al.*, 2001).

HS give birth to a single pup, which actively swims and dives in shallow waters within hours of birth and throughout the nursing period. Similar to other phocids, pups have a short nursing period lasting 24 days (Bowen *et al.*, 1992; Bowen *et al.*, 2001) resulting in a rapid postpartum growth associated with a large deposit of lipid reserves in the blubber layer. This foraging and lactation cycle differs somewhat from other phocids that fast during the entire nursing period (ranging 4-50 days). The foraging and nursing cycle observed in HS is similar to, though comparatively shorter than, those observed in otariids (Boness *et al.*, 1994; Bowen *et al.*, 2001). Bowen *et al.*, (1992) suggest the harbor seal lactation strategy is the result of smaller lipid reserves in the adult female compared to other phocids.

Given the high energy demand and rapid growth and development in both harbor seal and Steller sea lion juveniles and pups, I hypothesize that age or body size will be associated with circulating concentrations of hormones, leukocyte counts and *in vitro* proliferation of PBMC.

1.3 Body Condition

Disease and nutrition are considered the major external biotic factors influencing growth in vertebrates (Vander *et al.*, 1986). Body condition indices (BCI) have been

applied in livestock to assess health and production measures, e.g. sheep (Caldeira *et al.*, 2007), and to wild populations of terrestrial mammals, e.g. bears (Cattet *et al.*, 2002). Increasingly there has been an interest in quantifying body condition in pinnipeds as a measure of health and fitness (Castellini and Calkins, 1993; Arnould and Hindell, 1999; Jonker and Trites, 2000; Pitcher *et al.*, 2000; Trites and Jonker, 2000; Arnould and Warneke, 2002; Rea, 2002). BCI in pinnipeds have been proposed based on various combinations of mass, standard length, axillary girth, and blubber thickness (Pitcher *et al.*, 2000; Trites and Jonker, 2000; Rea, 2002). These calculated indices have been used for prediction of energy reserves, reproductive success and pup survival.

Body mass and condition are influenced by an individual's disease and nutritional state and therefore may act as a proxy for assessing an individual's or population's health. The potential impact of body condition on an animal's ability to survive would be greatest during times of high energy demands such as pregnancy, lactation and periods of development and rapid growth. Adult female otariids (Antarctic fur seals, *Arctocephalus gazella*) exhibited reduced pup production in years following an El Niño Southern Oscillation (Guinet *et al.*, 1994) and larger, older females gave birth to larger pups (Lunn and Boyd, 1993). In southern elephant seals (*Mirounga leonina*) larger females gave birth to larger pups (regardless of pup sex) and male pups were heavier than female pups at the time of birth (Arnbom *et al.*, 1994). Similarly, mass of HS at birth and postnatal growth rates were influenced by the mass of the adult female, with heavier females giving birth to heavier pups with a faster growth rate (Ellis *et al.*, 2000; Bowen *et al.*, 2001). Body mass and lipid stores in HS were correlated with survival during the first year of independence (Muelbert *et al.*, 2003; Harding *et al.*, 2005). Overall, there is strong evidence supporting the importance of body mass at birth or weaning as predictive of survivorship of pups in phocids and otariids (Baker and Fowler, 1992; Kjellquist *et al.*, 1995; Harding *et al.*, 2005). However, the relationships between BCI or mass with health parameters such as complete blood cell (CBC) count, immune function (e.g. PBMC proliferation), or circulating hormone concentrations have not been fully explored in pinnipeds.

Seasonally breeding mammals such as pinnipeds commonly exhibit robust annual cycles in body mass and energy demands (Lincoln and Richardson, 1998; Rousseau *et al.*, 2003). BCI or body mass and nutritional status showed seasonality in phocids (Ryg *et al.*, 1990; Beck *et al.*, 1993; Renouf *et al.*, 1993; Nilssen *et al.*, 2001; Chabot and Stenson, 2002) and otariids (Kumagai *et al.*, 2006; Rea *et al.*, 2007; Williams *et al.*, 2007; Jeanniard du Dot *et al.*, 2008). Changes in body mass and/or nutritional status are characterized by alterations in concentrations of many hormones that are related to body condition (Bhat *et al.*, 2003). Similar seasonality occurs in pinnipeds (Rosen and Renouf, 1997; Rosen and Renouf, 1998; Kumagai *et al.*, 2006; Mellish *et al.*, 2007; Rosen and Kumagai, 2008), which has been associated with changes in cortisol and thyroid hormones (Riviere *et al.*, 1977; Ashwell-Erickson *et al.*, 1986; Boily, 1996; Oki and Atkinson, 2004; Rosen and Kumagai, 2008; Jeanniard du Dot *et al.*, 2009). Specifically, decreases in lipid stores and mass were associated with an increase in circulating cortisol concentration and decrease in total T₄ and T₃ concentrations in SSL during experimental food restriction (Rosen and Kumagai, 2008; Jeanniard du Dot *et al.*, 2009). Many of these studies have focused on periods of high energy demands such as lactation, natural or experimental fasting, or molting. Previous studies assessing the relationship of body mass or condition with hormones in SSL primarily focused on juvenile and sub-adults (Mellish *et al.*, 2006; Jeanniard du Dot *et al.*, 2008; Rosen and Kumagai, 2008; Jeanniard du Dot *et al.*, 2009; Rea *et al.*, 2009) and did not assess the impact of mass on immune parameters. Conversely, studies on HS have generally focused on pups and the influence of body mass or age on either changes in hormone concentrations associated with weaning (Riviere *et al.*, 1977; Ashwell-Erickson *et al.*, 1986; Boily, 1996) or the postnatal development of the immune system by assessing leukocyte counts and *in vitro* proliferation of PBMC (Swart *et al.*, 1993; Ross *et al.*, 1994; Nielsen, 1995; Levin *et al.*, 2005; Kakuschke *et al.*, 2008a). Therefore, this study assesses how age or body size are associated with circulating concentrations of hormones as well as circulating total and differential leukocyte counts and the *in vitro* proliferation of PBMC in SSL pups. Further, this study seeks to build upon the abundance of research on hormone concentrations,

leukocyte counts and PBMC proliferation in HS by assessing how body mass influences the physiological response to an acute stressor (exogenous ACTH) on cortisol, aldosterone, thyroxine and triiodothyronine concentrations, total and differential leukocyte counts and *in vitro* PBMC proliferation in juvenile HS.

1.4 Endocrine System

1.4.1 Hypothalamic-Pituitary-Adrenal Axis

Hypothalamic-pituitary-adrenal (HPA) axis activity varies between and within species depending on life history traits, environmental conditions, breeding status, sex and age. The HPA axis and its endpoint products, cortisol, the prominent glucocorticoid (GC) hormone in pinnipeds (DeRoos and Bern, 1961; Sangalang and Freeman, 1976) and aldosterone, a mineralocorticoid, play key roles in physiology and behavior of animals. GC are synthesized in the zona fasciculata of the adrenal cortex following stimulation by ACTH and inhibit peripheral tissues such as skeletal muscle from utilizing glucose and enhance the mobilization of fat stores to non-neuronal tissues. In pinnipeds, cortisol plays a significant role during fasting and/or lactation (Engelhard *et al.*, 2002; Guinet *et al.*, 2004; Jeanniard du Dot *et al.*, 2009), molt (Riviere *et al.*, 1977; Ashwell-Erickson *et al.*, 1986), and potentially during diving (Zapol *et al.*, 1979).

Pinnipeds exhibit a typically mammalian structure in the adrenal gland with an increase in connective tissue and surface lobulation (Ridgway, 1972). Adrenal gland mass increased with age in HS with relative adrenal gland mass decreasing with body mass (Sucheston and Cannon, 1980; Bragulla *et al.*, 2004). The medullary area to cortex ratio has been described as low (Sucheston and Cannon, 1980). The cortex shows typical mammalian zonation with the of zona glomerulosa accounting for nearly half the cortical area, which is larger compared to terrestrial mammals (DeRoos and Bern, 1961; Sucheston and Cannon, 1980). The increase in area of the zona glomerulosa was attributed to the need to maintain electrolyte balance while living in a marine environment. Aldosterone is secreted from the cells of the zona glomerulosa and is regulated by the renin-angiotensin system in pinnipeds (Malvin *et al.*, 1975; Ortiz *et al.*,

2000; Houser *et al.*, 2001; Ortiz *et al.*, 2003a; Ortiz *et al.*, 2006). Secretion of aldosterone is stimulated by angiotensin II and induces the absorption of Na^+ in the distal tubule of the nephron maintaining water and electrolyte balance. Previous studies have also demonstrated that aldosterone is secreted in response to a stressor in mammals. For example, aldosterone concentrations increased during acute stress response (Thomson and Geraci, 1986; St. Aubin *et al.*, 1996; Gardiner and Hall, 1997; Contreras *et al.*, 2004; Romano *et al.*, 2004) and during exogenous ACTH challenges (St. Aubin and Geraci, 1986; Gulland *et al.*, 1999) in association with increased concentrations of cortisol, a well known stress hormone.

Stress has been defined as a state of threatened homeostasis resulting from internal or external environmental change (Stratakis and Chrousos, 1995; Leonard, 2005). Stress falls into one of two categories based on the duration or frequency of the stressor. Stress marked by a long duration or with frequent recurrence over an extended period is classified as chronic, while stressors having a sudden onset, sharp rise, and short course are classified as acute. Animals or populations in poor body condition may be experiencing chronic stress associated with a reduction in prey abundance or quality often referred to as nutritional stress (Merrick *et al.*, 1987; Calkins *et al.*, 1998; Anderson *et al.*, 1999; Trites and Donnelly, 2003). Capture and handling methods routinely used during blood collection in wild mammal populations is classified as an acute stressor and has been shown to result in elevated cortisol concentrations (Engelhard *et al.*, 2002; Romero *et al.*, 2008) and alter hematological parameters (Castellini *et al.*, 1996; Cattet *et al.*, 2003). Cortisol concentration (approximately 130 ng/ml) was elevated following capture in juvenile SSL compared to concentrations measured following acclimation to temporary captivity (74 ± 22 ng/ml) (Petrauskas *et al.*, 2008) and from captive juvenile and adult SSL (Mashburn and Atkinson, 2004; Rosen and Kumagai, 2008).

Elevated cortisol concentrations have been shown to suppress immune function including PBMC proliferation in piglets (Westly and Kelley, 1984) while having no influence in lambs (Minton and Blecha, 1990; Coppinger *et al.*, 1991) or calves (Manak, 1986). The effect of elevated GC on immune parameters is dependent on several

variables including the concentration of GC induced by an acute stressor, the immune parameter or function, and the species being studied (Anderson *et al.*, 1999; Bilandzic *et al.*, 2006). An acute stress response is generally considered adaptive while exposure to chronic stressors and the associated prolonged elevated concentrations of GC is detrimental to the health and survival of individuals. Morphological changes in adrenal gland size, medulla to cortex ratio, and individual zones have been associated with chronic stress and disease. Adrenal necrosis associated with herpes virus inclusions in harbor seal pups that died during rehabilitation showed marked lymphoid atrophy in the thymus, spleen and lymph nodes and significant lymphopenia (Gulland *et al.*, 1997; Gulland *et al.*, 1999), though the mortality was not attributed to adrenal insufficiency. The acute lymphoid depletion observed in several lymph nodes (tracheobronchial, axillary, mesenteric or jejuna) in leopard seals (*Hydrurga leptonyx*) was attributed to endogenous steroid release (Gray *et al.*, 2006). However, previous studies on the acute stress response in pinnipeds (St. Aubin and Geraci, 1986; Gulland *et al.*, 1999; Mashburn and Atkinson, 2008) have not assessed the impact of an acute stressor on the total and differential leukocyte count or *in vitro* proliferation of PBMC. Therefore, this study assessed the effect of exogenous ACTH on *in vitro* PBMC proliferation in addition to total and differential leukocyte counts in HS as a means of assessing the impact of an acute stressor.

1.4.2 Thyroid Gland and Hormones

Thyroid hormones including total thyroxine (T_4), free T_4 and total triiodothyronine (T_3) are essential for growth and development, metabolism and thermoregulation. Thyroid gland mass increased with age with sexually mature adult thyroid weighing about twice that of immature harbor seals (Harrison *et al.*, 1962); however, no difference in mass was observed between sexes in pups or adults (Harrison *et al.*, 1962; Little, 1991). Two periods of growth in the thyroid gland were suggested, the first period occurred during late gestation with the second occurring near sexual maturation coinciding with an increase in body mass. Thyroid hormones were also

affected by age in seals with pups having higher concentrations in T_4 and T_3 , which has been attributed to the high metabolic and thermoregulatory needs of pups (Hall *et al.*, 1998; Haulena *et al.*, 1998; Woldstad and Jenssen, 1999).

1.4.3 Leptin

Leptin is secreted by adipose tissue (Friedman and Halaas, 1998; Kershaw and Flier, 2004) and circulating concentrations have been correlated with fat stores and respond to changes in energy balance in terrestrial mammals (Friedman and Halaas, 1998). Specifically, in human and rodents there is a strong positive relationship between leptin concentrations and body fat mass (Ahima *et al.*, 1996). In horses (Buff *et al.*, 2002), cows and sheep (Chilliard *et al.*, 2001) body condition and fat mass were similarly related to serum leptin concentrations. Further, there is increasing evidence that leptin has systemic effects including regulation of neuroendocrine, immune function and development. Reduced circulating concentrations of leptin during malnutrition have been shown to directly influence acquired immunity by suppressing T lymphocyte proliferation (Lord *et al.*, 1998).

Leptin concentration has a clear relationship with fat mass in some mammals; however, the relationship has not been observed in other mammalian species. These findings have lead some researchers to suggest that carnivores lack a detectable relationship between body condition and leptin concentrations (Ahima and Flier, 2000; Nieminen *et al.*, 2001; Arnould *et al.*, 2002). For example, in terrestrial mammals, food restrictions leading to a loss in body fat mass was not associated with changes in leptin concentration in the raccoon dog (*Nyctereutes procyonoides*) (Nieminen *et al.*, 2004). Similarly, leptin concentrations in northern elephant seal (*Mirounga angustirostris*) (Ortiz *et al.*, 2001; Ortiz *et al.*, 2003b) and Antarctic fur seal pups (Arnould *et al.*, 2002) did not correlate to fat or body mass. However, adult elephant seals underwent a significant decrease in leptin associated with decreasing lipid stores while fasting during periods of reproduction and molt (Guilherme *et al.*, 2004). During periods of fasting, serum leptin concentrations were higher in adult elephant seals (Guilherme *et al.*, 2004)

compared to pups (Ortiz *et al.*, 2001; Ortiz *et al.*, 2003b). These findings may be reflective of the role of the blubber layer during fasting and postnatal development with the lower levels of leptin observed in otariid pups being reflective of the comparatively low lipid stores (Brandon *et al.*, 2005) and smaller blubber layer (Castellini *et al.*, 1993; Trites and Jonker, 2000) observed in otariid pups.

Fluctuating leptin concentrations not associated with changes in fat or body mass have been previously reported in pinnipeds (Arnould *et al.*, 2002; Mashburn and Atkinson, 2008). Mashburn and Atkinson (2008) reported a large amount of variability in serum leptin concentrations in response to an exogenous ACTH challenge in juvenile SSL. Difficulty in quantifying leptin in SSL (Rosen and Kumagai, 2008) and other pinnipeds (Arnould *et al.*, 2002; Ortiz *et al.*, 2003b) have been reported with low concentrations or no correlation to body mass or lipid stores, leading several researchers to propose that leptin is not a prominent signal of fat reserves in pinnipeds (Ortiz *et al.*, 2001; Arnould *et al.*, 2002; Rosen and Kumagai, 2008). Therefore, the relationship of leptin and body condition or mass remains unclear in pinnipeds. This study assessed leptin concentrations in SSL pups of known age and mass during the postnatal period to assess the relationship between mass and leptin concentrations during a period of rapid growth associated with increasing lipid stores in otariids.

1.5 Immune System

1.5.1 Immune Organs and Leukocytes

The vertebrate immune system has two types of responses, the innate (natural) and acquired (adaptive) responses. The innate response is the host defense that exists prior to exposure to a specific antigen and involves phagocytic cells (neutrophils, monocytes, and macrophages), inflammatory mediators (basophils and eosinophils) and natural killer cells. Neutrophils circulate in the peripheral blood for 7-10 hrs (e.g. human) following which they migrate into tissues where they will remain for a few days (Goldsby *et al.*, 2003). Monocytes circulate in the blood for about 8 hrs (e.g. human) and then migrate into tissues and differentiate into macrophages or dendritic cells (Goldsby *et al.*,

2003). During an infection, the number of circulating neutrophils increases (leukocytosis) and are generally the first cells to arrive at a site of inflammation. Similarly, circulating neutrophil counts increase rapidly with a concurrent decrease in lymphocyte and monocyte counts during an acute stress response in association with elevated GC concentrations (Dhabhar *et al.*, 1995; Dhabhar *et al.*, 1996; Anderson *et al.*, 1999).

Lymphocytes are responsible for the adaptive immune response in vertebrates with the majority of lymphocytes found in lymph. Lymphocytes mature and become committed to a particular antigenic lineage in the thymus (T cells) or in the bone marrow (B cells). Mature lymphocytes leave the primary lymphoid organs and circulate in the blood and lymphatic system. The adaptive response is generated in the spleen, lymph nodes, and the mucosa-associated lymphoid tissue leading to the proliferation of antigen-specific B and T lymphocytes.

The primary organs of the mammalian immune system are the thymus and bone marrow where lymphocytes mature. The thymus in pinnipeds has not been well described in the literature. Smodlaka *et al.*, (2009) noted that the thymus of one juvenile ringed seal (*Pusa hispida*) was bilobed and easily identified while only remnants of the thymus were present in an adult. Involution of the thymus associated with aging has been described in several species including the dog (Newman, 1971), dolphin (Clark *et al.*, 2005), cattle (Lubis *et al.*, 1982) and human (Suster and Rosai, 1990). It is therefore likely that the thymus in pinnipeds undergoes involution following the onset of sexual maturity as in terrestrial mammals.

The secondary lymphoid organs include skin, lymph nodes, and the spleen. Though often overlooked, the first barrier to infection is the skin and the mucosal membranes of an animal. Histology of skin in several species of pinnipeds showed increased thickness of the epidermis and numerous sebaceous glands (Montagna and Harrison, 1957). The integument of several pinniped species, including the harbor seal were positive for the presence of β -glucan receptors in association with skin glands, outer epithelial sheath of the hair follicles and in the epidermis, supporting a role of pinniped skin in the innate immune system (Meyer *et al.*, 2008).

The spleen of pinnipeds appears to have several functions including immunologic, haematopoiesis and blood storage (Schumacher and Welsch, 1987). The spleen has been described as moderately large (Slijper, 1958) and increases in mass with body mass (Scheffer, 1960) but accounts for approximately 0.11-0.40 % of the total body weight in Cape fur seal (Otariid; *Arctocephalus pusillus*) (Stewardson *et al.*, 1999) and the northern fur seal (Otariid; *Callorhinus ursinus*) (Scheffer, 1960), which is comparable to terrestrial mammals. The spleen is enclosed in a thick capsule with prominent trabecule and innervations of smooth muscle cells (Schumacher and Welsch, 1987; Gray *et al.*, 2006). Stewardson *et al.*, (1999) determined that the spleen, along with the heart and liver, in the Cape fur seal grew at a faster rate than the body and other organs. The spleen grew approximately 33% per year until at least 10 years of age with varying degrees of development in the red and white pulp of the parenchyma, which may indicate ontogeny changes, disease status, or both. The noted size and growth of the spleen has generally been attributed to the spleen's role in sequestering red blood cells and diving ability of these species (Cabanac *et al.*, 1989; Ponganis *et al.*, 1997). However, the spleen in pinnipeds (both phocids and otariid) has a prominent lymphatic part with well developed follicles and numerous plasma cells in the white pulp and eosinophils in the red pulp. The presence of eosinophils in the spleen was attributed to the relatively high load of gastrointestinal parasites (Schumacher and Welsch, 1987; Stewardson *et al.*, 1999; Gray *et al.*, 2006). Eosinophils are motile phagocytic cells that can migrate from the blood into the tissue spaces and are thought to play a role in the defense against parasitic organisms.

The lymph nodes of pinnipeds exhibit the general mammalian pattern both structurally and histologically with lymphoid follicles located within the cortex (Welsch *et al.*, 1997; Gray *et al.*, 2006). In both phocids and otariids, the capsule and trabecular penetrating the parenchyma of lymph nodes are broad containing smooth muscle cells within a network of collagenous and elastic fibers. Similar structures have been found in odontoceti (toothed whales) and have been proposed to enable active movement and filtration of lymph fluid by capsular contraction (Romano *et al.*, 1993; Cowan and Smith, 1999). The degree of development of lymphoid tissue was variable between individuals

(Stewardson *et al.*, 1999). Kuttyrev *et al.*, (2008) found ontogenetic changes in the mesenteric lymph nodes of the Baikal Seal (*Pusa sibirica*) including a dramatic increase in the capsule and trabeculae with age. These changes were associated with a decrease in the relative area of the cortex, secondary lymph follicles, paracortical zone and medulla resulting in an overall decrease of the corticomedullary index with age.

The mesenteric lymph nodes of the harbor and grey (*Halichoerus grypus*) seals exhibited numerous fully developed large secondary follicles and broad T cell regions with most follicles appearing as germinal centers indicative of an immunological response to an antigenic challenge (Welsch *et al.*, 1997). These mesenteric lymph nodes were collected from healthy individuals lacking any gross signs of disease and these observations were attributed to the exposure to antigens within the coastal marine or terrestrial environment.

Lymph nodes obtained during epidemics of harbor seal (Schumacher *et al.*, 1990) or from compromised leopard seals (Gray *et al.*, 2006) were lymphocyte depleted compared to other mammals or healthy animals of the same species (Stewardson *et al.*, 1999). The observed lymphocyte depletion was attributed to either the natural disease process or the result of endogenous steroid release associated with restraint and sedation (Gray *et al.*, 2006). Similar to findings in the spleen, eosinophils were noted in the lymph nodes in both phocids (Boily, 1996) and otariids (Simpson and Gardner, 1972; Welsch *et al.*, 1997) and were attributed to the prevalence of parasites in these species.

1.5.2 Hematology and PBMC Proliferation

Hematology has been routinely used to assess the health of individuals and populations of pinnipeds including SSL (Castellini *et al.*, 1993; Bishop and Morado, 1995; Rea *et al.*, 1998) and HS (McConnell and Vaughan, 1983; Ross *et al.*, 1993; Morgan *et al.*, 1998; Lander *et al.*, 2003; Hasselmeier *et al.*, 2008). Further, some hematological parameters such as hematocrit (Hct), hemoglobin (Hb), white blood cell (WBC) and platelet count (Plt) have been used for assessing body condition and nutritional status or stress in several species of pinnipeds (Thompson *et al.*, 1997; Rosen

et al., 2004; Trumble *et al.*, 2006). Across mammalian species and in both sexes the number of circulating total WBC count increased with body mass in adults (Nunn *et al.*, 2003; Nunn *et al.*, 2009). This increase was driven predominantly by circulating neutrophil and monocyte counts with a concurrent decrease in lymphocyte counts. Significant differences in total and differential WBC counts were found between immature and mature mammals across taxa including carnivores (Nunn *et al.*, 2009). The strongest difference was found in circulating lymphocyte counts with immature individuals having higher lymphocyte counts than both mature males and females of the same species. Similar observations have been reported in pinnipeds. Nielsen (1995) found neutrophils (%) increased while lymphocytes (%) decreased with age (1.5-17.5 yrs) in HS. The decrease in lymphocytes (%) was associated with a decrease in response to *in vitro* mitogen stimulation of PBMC. However, Kakuschke *et al.*, (2008b) reported higher PBMC proliferation in HS pups compared to adults. Taken together these findings are suggestive that the age-related changes in PBMC proliferation are not linear with age but that pups undergo an initial increase prior to a decrease in proliferation of PBMC in adults. Similar studies have not been done on SSL. Mellish *et al.*, (2006) sampled SSL pups (6–10 months) and juveniles (1–3 years) concluding that pups had significantly lower WBC counts than juveniles; however, differential leukocyte counts and PBMC proliferation were not included in the study. No study has assessed the age-related change in total and differential leukocyte counts or PBMC proliferation in SSL less than 6 months of age. Therefore, this project sampled SSL pups during the early postnatal period to assess changes in total and differential leukocyte counts and proliferation of PBMC as stimulated by concanavalin A (ConA) and lipopolysaccharide (LPS).

1.6 Literature Cited

- Agarwal, S. K., Marshall Jr., G. D., 1998. Glucocorticoid-induced type1/type2 cytokine alterations in humans: A model for stress-related immune dysfunction. *J. Interferon Cytokine Res.* 18, 1059-1068.
- Ahima, R. S., Prabakaran, D., Mantzoros, C., Qu, D., Lowll, B., Maratos-Filter, E., Flier, J. S., 1996. Role of leptin in the neuroendocrine response to fasting. *Nature.* 382, 250-252.
- Ahima, R. S., Flier, J. S., 2000. Leptin. *Ann. Rev. Physiol.* 62, 413-437.
- Anderson, B. H., Watson, D. L., Colditz, I. G., 1999. The effect of dexamethasone on some immunological parameters in cattle. *Vet. Res. Commun.* 23, 399-413.
- Arnbom, T., Fedak, M. A., Rothery, P., 1994. Offspring sex ratio in relation to female size in southern elephant seals, *Mirounga leonina*. *Behav. Ecol. Sociobiol.* 35, 373-378.
- Arnould, J. P. Y., Boyd, I. L., Socha, D. G., 1996. Milk consumption and growth efficiency in Antarctic fur seal (*Arctocephalus gazella*) pups. *Can. J. Zool.* 74, 254-266.
- Arnould, J. P. Y., Hindell, M. A., 1999. The composition of Australian fur seal (*Arctocephalus pusillus doriferus*) milk throughout lactation. *Physiol. Biochem. Zool.* 72, 605-612.
- Arnould, J. P. Y., Morris, M. J., Rawlins, D. R., Boyd, I. L., 2002. Variation in plasma leptin levels in response to fasting in Antarctic fur seals (*Arctocephalus gazella*). *J. Comp. Physiol. B.* 172, 27-34.
- Arnould, J. P. Y., Warneke, R. M., 2002. Growth and condition in Australian fur seals (*Arctocephalus pusillus doriferus*) (Carnivora: Pinnipedia). *Aust. J. Zool.* 50, 53-66.
- Ashwell-Erickson, S., Fay, F. H., Elsner, R., 1986. Metabolic and hormonal correlates of molting and regeneration of pelage in Alaskan harbor and spotted seals (*Phoca vitulina* and *Phoca largha*). *Can. J. Zool.* 64, 1086-1094.
- Atkinson, S., DeMaster, D. P., Calkins, D. G., 2008. Anthropogenic causes of the western Steller sea lion *Eumetopias jubatus* population decline and their threat to recovery. *Mamm. Rev.* 38, 1-18.

- Baker, A. R., Loughlin, T. R., Burkanov, V., Matson, C. W., Trujillo, R. G., Calkins, D. G., Wickliffe, J. K., Bickham, J. W., 2005. Variation of mitochondrial control region sequences of Steller sea lions: The three-stock hypothesis. *J. Mammal.* 86, 1075-1084.
- Baker, J. D., Fowler, C. W., 1992. Pup weight and survival of northern fur seals *Callorhinus ursinus*. *J. Zool. Lond.* 227, 231-238.
- Beck, G. G., Smith, T. G., Hammill, M. O., 1993. Evaluation of body condition in the Northwest Atlantic harp seal (*Phoca groenlandica*). *Can. J. Fish. Aquat. Sci.* 50, 1372-1381.
- Berta, A., Adam, P.J., 2001. Evolutionary biology of pinnipeds, In: Mazin, J.-M., Buffrenil, V.D. (Eds.) *Secondary Adaptation of Tetrapods to Life in Water*. Verlag Dr. Friedrich Pfeil. pp. 235-260.
- Bhat, G. K., Hamm, M. L., Igiertsem, J. U., Mann, D. R., 2003. Does leptin mediate the effect of photoperiod on immune function in mice? *Biol. Reprod.* 69, 30-36.
- Bickham, J. W., Patton, J. C., Loughlin, T. R., 1996. High variability for control-region sequences in a marine mammal: Implications for conservation and biogeography of Steller sea lions (*Eumetopias jubatus*). *J. Mammal.* 77, 95-108.
- Bickham, J. W., Loughlin, T. R., Calkins, D. G., Wickliffe, J. K., Patton, J. C., 1998. Genetic variability and population decline in Steller sea lions from the Gulf of Alaska. *J. Mammal.* 79, 1390-1395.
- Bilandzic, N., Zuric, M., Lojkic, M., Simic, B., Milic, D., Bara, I., 2006. Cortisol and immune measures in boars exposed to three-day administration of exogenous adrenocorticotrophic hormone. *Vet. Res. Commun.* 30, 433-444.
- Bishop, D. H., Morado, J. F., 1995. Results on blood cell morphology and differential blood cell counts from seventeen Steller sea lion *Eumetopias jubatus* pups. *Dis. Aquat. Org.* 23, 1-6.
- Boily, P., 1996. Metabolic and hormonal changes during the molt of captive gray seals (*Halichoerus grypus*). *Am. J. Physiol. Regul. Integr. Comp. Physiol.* 270, 1051-1058.
- Boness, D. J., Bowen, W. D., Oftedal, O. T., 1994. Evidence of a maternal foraging cycle resembling that of otariid seals in a small phocid, the harbor seal. *Behav. Ecol. Sociobiol.* 34, 95-104.

- Boness, D. J., Bowen, W. D., 1996. The evolution of maternal care in pinnipeds. *BioScience*. 46, 645-654.
- Bonner, W. N., 1984. Lactation strategies in pinnipeds: Problems for a marine mammalian group. *Symp. Zool. Soc. Lond.* 51, 253-272.
- Bowen, W. D., 1991. Behavioural ecology of pinniped neonates. In: Renouf, D. (ed) *The behavioural ecology of pinniped*. Chapman and Hall, p. 66-127.
- Bowen, W. D., Oftedal, O. T., Boness, D. J., 1992. Mass and energy transfer during lactation in a small phocid, the harbor seal (*Phoca vitulina*). *Physiol. Zool.* 65, 844-866.
- Bowen, W. D., Ellis, S. L., Iverson, S. J., Boness, D. J., 2001. Maternal effects on offspring growth rate and weaning mass in harbour seals. *Can. J. Zool.* 79, 1088-1101.
- Bowen, W. D., Iverson, S. J., McMillan, J. I., Boness, D. J., 2006. Reproductive performance in grey seals: age-related improvement and senescence in a capital breeder. *J. Anim. Ecol.* 75, 1340-1351.
- Boyd, I. L., 1998. Time and energy constraints in pinniped lactation. *Am. Nat.* 152, 717-728.
- Bragulla, H., Hirschberg, R. M., Schlotfeldt, U., Stede, M., Budras, K.-D., 2004. On the structure of the adrenal gland of the common seal (*Phoca vitulina vitulina*). *Anat. Histol. Embryol.* 33, 263-272.
- Brandon, E. A. A., Calkins, D. G., Davis, R. W., 2005. Neonatal growth of Steller sea lion (*Eumetopias jubatus*) pups in Alaska. *Fish. Bull.* 103, 246-257.
- Buff, P. R., Dodds, A. C., Morrison, C. D., Whitley, N. C., McFadin, E. L., Daniel, J. A., Djiane, J., Keisler, D. H., 2002. Leptin in horses: tissue localization and relationship between peripheral concentrations of leptin and body condition. *J. Anim. Sci.* 80, 2942-2948.
- Burek, K. A., Gulland, F. M. D., Sheffield, G., Keyes, E., Spraker, T. R., Smith, A. W., Skilling, D. E., Evermann, J., Stott, J. L., Trites, A. W., 2003. Disease agents in Steller sea lions in Alaska: a review and analysis of serology data from 1975-2000. In *Fisheries Centre Research Reports* (Vancouver, University of British Columbia), p. 26.

- Burek, K. A., Gulland, F. M. D., Sheffield, G., Beckmen, K., Keyes, E., Spraker, T., Smith, A. W., Skilling, D. E., Evermann, J. F., Stott, J. L., Saliki, J. T., Trites, A. W., 2005. Infectious disease and the decline of Steller sea lions (*Eumetopias jubatus*) in Alaska, USA: insights from serologic data. *J. Wildl. Dis.* 41, 512-524.
- Burek, K. A., Gulland, F. M. D., O'Hara, T. M., 2008. Effects of climate change on arctic marine mammal health. *Ecol. Appl.* 18, S126-S134.
- Cabanac, A., Folkow, L. P., Blilx, A. S., 1989. Volume capacity and contraction control of the seal spleen. *J. Appl. Physiol.* 82, 1989-1994.
- Caldeira, R. M., Belo, A. T., Santos, C. C., Vazques, M. I., Portugal, A. V., 2007. The effect of body condition score on blood metabolites and hormonal profiles in ewes. *Sm. Rumin. Res.* 68, 233-241.
- Calkins, D. G., Becker, E. F., Pitcher, K. W., 1998. Reduced body size of female Steller sea lions from a declining population in the Gulf of Alaska. *Mar. Mamm. Sci.* 14, 232-244.
- Castellini, J. M., Melselman, H. J., Castellini, M. A., 1996. Understanding and interpreting hematocrit measurements in pinnipeds. *Mar. Mamm. Sci.* 12, 251-264.
- Castellini, M. A., Calkins, D. G., 1993. Mass estimates using body morphology in Steller sea lions. *Mar. Mamm. Sci.* 9, 48-54.
- Castellini, M. A., Davis, R. W., Loughlin, T. R., Williams, T. M., 1993. Blood chemistries and body condition of Steller sea lion pups at Marmot Island, Alaska. *Mar. Mamm. Sci.* 9, 202-208.
- Cattet, M. R. L., Caulkett, N. A., Obbard, M. E., Stenhouse, G. B., 2002. A body-condition index for ursids. *Can. J. Zool.* 80, 1156-1161.
- Cattet, M. R. L., Christison, K., Caulkett, N. A., Stenhouse, G. B., 2003. Physiologic responses of grizzly bears to different methods of capture. *J. Wildl. Dis.* 39, 649-654.
- Chabot, D., Stenson, G. B., 2002. Growth and seasonal fluctuations in size and condition of male Northwest Atlantic harp seals *Phoca groenlandica*: an analysis using sequential growth curves. *Mar. Ecol. Prog. Ser.* 227, 25-42.

- Chilliard, Y., Bonnet, M., Delavaud, C., Faulconnier, Y., Leroux, C., Djiane, J., Bocquier, F., 2001. Leptin in ruminants. Gene expression in adipose tissue and mammary gland, and regulation of plasma concentration. *Domest. Anim. Endocrinol.* 21, 271-295.
- Clark, C. A., Burns, J. M., Schreer, J. F., Hammill, M. O., 2007. A longitudinal and cross-sectional analysis of total body oxygen store development in nursing harbor seals (*Phoca vitulina*). *J. Comp. Physiol. B.* 177, 217-227.
- Clark, L. S., Turner, J. P., Cowan, D. F., 2005. Involution of lymphoid organs in bottlenose dolphins (*Tursiops truncatus*) from the Western Gulf of Mexico: Implications for life in an aquatic environment. *Anat. Rec. A.* 282A, 67-73.
- Contreras, L. N., Arregger, A. L., Persi, G. G., Gonzalez, N. S., Cardoso, E. M., 2004. A new less-invasive and more informative low-dose ACTH test: salivary steroids in response to intramuscular corticotrophin. *Clin. Endocrinol.* 61, 675-682.
- Coppinger, T. R., Minton, J. E., Reddy, P. G., Blecha, F., 1991. Repeated restraint and isolation stress in lambs increases pituitary-adrenal secretions and reduces cell-mediated immunity. *J. Anim. Sci.* 69, 2808-2814.
- Costa, D. P., 1991. Reproductive and foraging energetic of pinnipeds: implications for life-history patters. In: D. Renouf, (Ed.), *Behavior of Pinnipeds*. Chapman & Hall, London, p. 300-344.
- Cowan, D. F., Smith, T. L., 1999. Morphology of the lymphoid organs of the bottlenose dolphin, *Tursiops truncatus*. *J. Anat.* 194, 505-517.
- DeMaster, D., P., Trites, A. W., Clapham, P., Mizroch, S. A., Wade, P., Small, R. J., Hoef, J. V., 2006. The sequential megafaunal collapse hypothesis: testing with existing data. *Prog. Oceanogr.* 68, 329-342.
- DeRoos, C. C., Bern, H. A., 1961. The corticoids of the adrenal of the California sea lion (*Zalophus californianus*). *Gen. Comp. Endocrinol.* 1, 275-285.
- Dhabhar, F. S., Miller, A. H., McEwen, B. S., Spencer, R. L., 1995. Effects of stress on immune cell distribution. *J. Immunol.* 154, 5511-5527.
- Dhabhar, F. S., Miller, A. H., McEwen, B. S., Spencer, R. L., 1996. Stress-induced changes in blood leukocytes distribution: role of adrenal steroid hormones. *J. Immunol.* 157, 1638-1644.

- Ellis, S. L., Bowen, W. D., Boness, D. J., Iverson, S. J., 2000. Maternal effects on offspring mass and stage of development at birth in the harbor seal, *Phoca vitulina*. J. Mammal. 81, 1143-1156.
- Engelhard, G. H., Brasseur, S. M. J. M., Hall, A. J., Burton, H. R., Reijnders, P. J. H., 2002. Adrenocortical responsiveness in southern elephant seal mothers and pups during lactation and the effect of scientific handling. J. Comp. Physiol. B. 172, 315-328.
- Friedman, J. M., Halaas, J. L., 1998. Leptin and the regulation of body weight in mammals. Nature. 395, 763-770.
- Fritz, L. W., Hinckley, S., 2005. A critical review of the regime shift-"junk food"-nutritional stress hypothesis for the decline of the western stock of Steller sea lion. Mar. Mamm. Sci. 21, 476-518.
- Frost, K. J., Lowry, L. F., Hoef, J. M. V., 1999. Monitoring the trend of harbor seals in Prince William Sound, Alaska, after the Exxon Valdez oil spill. Mar. Mamm. Sci. 15, 494-506.
- Frost, K. J., Simpkins, M. A., Small, R. J., Lowry, L. F., 2006. Development of diving by harbor seal pups in two regions of Alaska: use of the water column. Mar. Mamm. Sci. 22, 617-643.
- Gardiner, K. J., Hall, A. J., 1997. Diel and annual variation in plasma cortisol concentrations among wild and captive harbor seals (*Phoca vitulina*). Can. J. Zool. 75, 1773-1780.
- Goldsby, R. A., Kindt, T. J., Osborne, B. A., Kuby, J., 2003. Immunol. W.H. Freeman and Company, New York.
- Goldstein, T., Stephens, C. A., Jang, S. S., Conrad, P. A., Field, C., Dunn, J. L., Mellish, J.-A. E., 2007. Longitudinal health and disease monitoring in juvenile Steller sea lions (*Eumetopias jubatus*) in temporary captivity in Alaska compared with a free-ranging cohort. Aquat. Mamm. 33, 337-348.
- Gray, R., Canfield, P., Rogers, T., 2006. Histology of selected tissues of the leopard seal and implications for functional adaptations to an aquatic lifestyle. J. Anat. 209, 179-199.
- Guenette, S., Heymans, S. J. J., Christensen, V., Trites, A. W., 2006. Ecosystem models show combined effects of fishing, predation, competition, and ocean productivity on Steller sea lions (*Eumetopias jubatus*) in Alaska. Can. J. Fish. Aquat. Sci. 63, 2495-2517.

- Guilherme, C., Bianchini, A., Martinez, P. E., Robaldo, R. B., Colares, E. P., 2004. Serum leptin concentration during the terrestrial phase of the Southern elephant seal *Mirounga leonina* (Carnivora: Phocidae). *Gen. Comp. Endocrinol.* 139, 137-142.
- Guinet, C., Jouventin, P., Georges, J.-Y., 1994. Long term population changes of fur seals *Arctocephalus gazella* and *Arctocephalus tropicalis* on subantarctic (Crozet) and subtropical (St. Paul and Amsterdam) Islands and their possible relationship to El Nino Southern Oscillation. *Antarct. Sci.* 6, 473-478.
- Guinet, C., Servera, N., Mangin, S., Georges, J.-Y., Lacroix, A., 2004. Change in plasma cortisol and metabolites during the attendance period ashore in fasting lactating subantarctic fur seals. *Comp. Biochem. Physiol. A.* 137, 523-531.
- Gulland, F. M. D., Lowenstine, L. J., Lapointe, J. M., Spraker, T., King, D. P., 1997. Herpesvirus infection in stranded pacific harbor seals of coastal California. *J. Wildl. Dis.* 33, 450-458.
- Gulland, F. M. D., Haulena, M., Lowenstine, L. J., Munro, C., Graham, P. A., Bauman, J., Harvey, J., 1999. Adrenal function in wild and rehabilitated pacific harbor seals (*Phoca vitulina richardii*) and in seals with phocine herpesvirus-associated adrenal necrosis. *Mar. Mamm. Sci.* 15, 810-827.
- Haddad, J. J., Saade, N. E., Safieh-Garabedian, B., 2002. Cytokines and neuro-immune-endocrine interactions: a role for the hypothalamic-pituitary-adrenal revolving axis. *J. Neuroimmunol.* 133, 1-19.
- Hall, A. J., Green, N. J. L., Jones, K. C., Pomeroy, P. P., Harwood, J., 1998. Thyroid hormones as biomarkers in grey seals. *Mar. Poll. Bull.* 36, 424-428.
- Harding, K. C., Fugiwara, M., Axberg, Y., Harkonen, T., 2005. Mass-dependent energetics and survival in harbour seal pups. *Funct. Ecol.* 19, 129-135.
- Harrison, R. J., Rowlands, I. W., Whitting, H. W., Young, B. A., 1962. Growth and structure of the thyroid gland in the common seal (*Phoca vitulina*). *J. Anat.* 96, 3-16.
- Harvey, J. T., Brown, R. F., Mate, B. R., 1990. Abundance and distribution of harbor seals (*Phoca vitulina*) in Oregon, 1975-1982. *Northwest. Nat.* 71, 65-71.
- Hasselmeier, I., Fonfara, S., Driver, J., Siebert, U., 2008. Differential hematology profiles of free-ranging, rehabilitated, and captive harbor seals (*Phoca vitulina*) of the German North Sea. *Aquat. Mamm.* 34, 149-156.

- Haulena, M., St. Aubin, D. J., Duignan, P. J., 1998. Thyroid hormone dynamics during the nursing period in harbour seals, *Phoca vitulina*. Can. J. Zool. 76, 48-55.
- Holmes, E. E., York, A. E., 2003. Using age structure to detect impacts on threatened populations: a case study with Steller sea lions. Conserv. Biol. 17, 1794-1806.
- Holmes, E. E., Fritz, L. W., York, A. E., Sweeney, K., 2007. Age-structured modeling reveals long-term declines in the natality of western Steller sea lions. Ecol. Appl. 17, 2214-2232.
- Hood, W. R., Ono, K. A., 1997. Variation in maternal attendance patterns and pup behaviour in a declining population of Steller sea lions (*Eumetopias jubatus*). Can. J. Zool. 75, 1241-1246.
- Hoover-Miller, A., Atkinson, S., Conlon, S., Prewitt, J., Armato, P., *in press*. Persistent decline in abundance of harbor seals *Phoca vitulina richardsi* over three decades in Aialik Bay, an Alaskan tidewater glacial fjord. Mar. Ecol. Prog. Ser.
- Houser, D. S., Crocker, D. E., Webb, P. M., Costa, D. P., 2001. Renal function in suckling and fasting pups of the northern elephant seal. Comp. Biochem. Physiol. A: Mol. Integr. Physiol. 129, 405-415.
- Jeanniard du Dot, T., Rosen, D. A. S., Trites, A. W., 2008. Steller sea lions show diet-dependent changes in body composition during nutritional stress and recover more easily from mass loss in winter than in summer. J. Exp. Mar. Biol. Ecol. 367, 1-10.
- Jeanniard du Dot, T., Rosen, D. A. S., Richmond, J. P., Kitaysky, A. S., Zinn, S. A., Trites, A. W., 2009. Changes in glucocorticoid, IGF-I and thyroid hormones as indicators of nutritional stress and subsequent refeeding in Steller sea lions (*Eumetopias jubatus*). Comp. Biochem. Physiol. A. 152, 524-534.
- Jemison, L. A., Pendleton, G. W., Wilson, C. A., Small, R. J., 2006. Long-term trends in harbor seal numbers at Tugidak Island and Nanvak Bay, Alaska. Mar. Mamm. Sci. 22, 339-360.
- Jonker, R. A. H., Trites, A. W., 2000. The reliability of skinfold-calipers for measuring blubber thickness of Steller sea lion pups (*Eumetopias jubatus*). Mar. Mamm. Sci. 16, 757-766.
- Kakuschke, A., Valentine-Thon, E., Fonfara, S., Griesel, S., Rosenberger, T., Siebert, U., Prange, A., 2008a. Metal-induced impairment of the cellular immunity of newborn harbor seals (*Phoca vitulina*). Arch Environ. Contam. Toxicol. 55, 129-136.

- Kakuschke, A., Valentine-Thon, E., Griesel, S., Rosenberger, T., Mundry, R., Siebert, U., Prange, A., 2008b. Blood metal levels and metal-influenced immune functions of harbour seals in captivity. *Mar. Poll. Bull.* 56, 764-769.
- Kershaw, E. E., Flier, J. S., 2004. Adipose tissue as an endocrine organ. *J. Clin. Endocrinol. Metab.* 89, 2548-2556.
- Kjellqwist, S. A., Haug, T., Oritsland, T., 1995. Trends in age-composition, growth and reproductive parameters of Barents Sea harp seals, *Phoca groenlandica*. *ICES J. Mar. Sci.* 52, 197-208.
- Kovacs, K. M., Lavigne, D. M., 1986. Maternal investment and neonatal growth in phocid seals. *J. Anim. Ecol.* 55, 1035-1051.
- Kovacs, K. M., Lavigne, D. M., 1992. Maternal investment in otariid seals and walruses. *Can. J. Zool.* 70, 1953-1964.
- Kumagai, S., Rosen, D. A. S., Trites, A. W., 2006. Body mass and composition responses to short-term low energy intake are seasonally dependent in Steller sea lions (*Eumetopias jubatus*). *J. Comp. Physiol. B.* 176, 589-598.
- Kutyrev, I. A., Lamazhapova, G. P., Zhamsaranova, S. D., 2008. Structural organization of mesenteric lymph nodes in postnatal development of the Baikal seal, *Pusa sibirica* Gmel. *Biol. Bull.* 35, 389-393.
- Lander, M. E., Harvey, J. T., Gulland, F. M. D., 2003. Hematology and serum chemistry comparisons between free-ranging and rehabilitated harbor seal (*Phoca vitulina richardsi*) pups. *J. Wildl. Dis.* 39, 600-609.
- Le Boeuf, B. J., Crocker, D. E., 2005. Ocean climate and seal condition. *BMC Biol.* 3, 1-10.
- Leonard, B. E., 2005. The HPA and immune axes in stress: the involvement of the serotonergic system. *Eur. Psychiatr.* 20, S302-S306.
- Levin, M., De Guise, S. D., Ross, P. S., 2005. Association between lymphocyte proliferation and polychlorinated biphenyls in free-ranging harbor seal (*Phoca vitulina*) pups from British Columbia, Canada. *Environ. Toxicol. Chem.* 24, 1247-1252.
- Lincoln, G. A., Richardson, M., 1998. Photo-neuroendocrine control of seasonal cycles in body weight, pelage growth and reproduction: lessons from the HPD sheep model. *Comp. Biochem. Physiol. C.* 119, 283-294.

- Little, G. J., 1991. Thyroid morphology and function and its role in thermoregulation in the newborn southern elephant seal (*Mirounga leonina*) at Macquarie Island. *J. Anat.* 176, 55-69.
- Lord, G. M., Matarese, G., Howard, J. K., Baker, R. J., Bloom, S. R., Lechler, R. I., 1998. Leptin modulates the T-cell immune response and reverses starvation-induced immunosuppression. *Nature*. 394, 897-901.
- Loughlin, T. R., Rugh, D. J., Fiscus, C. H., 1984. Northern sea lion distribution and abundance: 1956-80. *J. Wildl. Manage.* 48, 729-740.
- Loughlin, T. R., 1998. The Steller sea lion: A declining species. *Biosph. Conserv.* 1, 91-98.
- Lubis, I., Ladds, P. W., Reilly, L. R., 1982. Age associated morphological changes in the lymphoid system of tropical cattle. *Res. Vet. Sci.* 32, 270-277.
- Lunn, N. J., Boyd, I. L., 1993. Effects of maternal age and condition on parturition and the perinatal period of Antarctic fur seals. *J. Zool. Lond.* 229, 55-67.
- Malvin, R. L., Ridgway, S. H., Cornell, L., 1975. Renin and aldosterone levels in dolphins and sea lions (40117). *Proc. Soc. Exp. Biol. Med.* 157, 665-668.
- Manak, R. C., 1986. Neonatal calf serum immunoregulation of lymphocyte blastogenic responses. *J. Anim. Sci.* 62, 759-771.
- Maniscalco, J. M., Parker, P., Atkinson, S., 2006. Interseasonal and interannual measures of maternal care among individual Steller sea lions (*Eumetopias jubatus*). *J. Mammal.* 87, 304-311.
- Markussen, N. H., Bjorge, A., Oritsland, N. A., 1989. Growth in harbour seals (*Phoca vitulina*) on the Norwegian coast. *J. Zool. Lond.* 219, 443-440.
- Mashburn, K. L., Atkinson, S., 2004. Evaluation of adrenal function in serum and feces of Steller sea lions (*Eumetopias jubatus*): influences of molt, gender, sample storage, and age on glucocorticoid metabolism. *Gen. Comp. Endocrinol.* 136, 371-381.
- Mashburn, K. L., Atkinson, S., 2008. Variability in leptin and adrenal response in juvenile Steller sea lions (*Eumetopias jubatus*) to adrenocorticotrophic hormone (ACTH) in different seasons. *Gen. Comp. Endocrinol.* 155, 352-258.

- Mathews, E. A., Pendleton, G. W., 2006. Declines in harbor seal (*Phoca vitulina*) numbers in galcier bay national park, Alaska, 1992-2002. *Mar. Mamm. Sci.* 22, 167-189.
- McConnell, L. C., Vaughan, R. W., 1983. Some blood values in captive and free-living common seals (*Phoca vitulina*). *Aquat. Mamm.* 10, 9-13.
- Mellish, J.-A. E., Calkins, D. G., Christen, D. R., Horning, M., Rea, L. D., Atkinson, S. K., 2006. Temporary captivity as a research tool: comprehensive study of wild pinnipeds under controlled conditions. *Aquat. Mamm.* 32, 58-65.
- Mellish, J.-A. E., Horning, M., York, A. E., 2007. Seasonal and spatial blubber depth changes in captive harbor seals (*Phoca vitulina*) and Steller's sea lions (*Eumetopias jubatus*). *J. Mammol.* 88, 408-414.
- Mellish, J.-A. E., Iverson, S. J., Bowen, W. D., Hammill, M. O., 1999. Fat transfer and energetics during lactation in the hooded seal: the roles of tissue lipoprotein lipase in milk fat secretion and pup blubber deposition. *J. Comp. Physiol. B.* 169, 377-390.
- Merrick, R. L., Loughlin, T. R., Calkins, D. G., 1987. Decline in abundance of the Northern sea lion, *Eumetopias jubatus*, in Alaska, 1956-1986. *Fish. Bull.* 85, 351-365.
- Meyer, W., Kloepper, J. E., Fleischer, L.-G., 2008. Demonstration of β -glucan receptors in the skin of aquatic mammals- a preliminary report. *Eur. J. Wildl. Res.* 54, 479-486.
- Millar, J. S., 1977. Adaptive features of mammalian reproduction. *Evol.* 31, 370-386.
- Minton, J. E., Blecha, F., 1990. Effect of acute stressors on endocrinological and immunological functions in lambs. *J. Anim. Sci.* 68, 3145-3151.
- Montagna, W., Harrison, R. J., 1957. Specialization in the skin of the seal (*Phoca vitulina*). *Am. J. Anat.* 100, 81-113.
- Morgan, L., Kumaresan, S., Thomas, C., MacWilliams, P., 1998. Hematology and chemistry reference values for free-ranging harbor seals (*Phoca vitulina*) and the effects of hemolysis on chemistry values of captive harbor seals. *J. Zoo Wildl. Med.* 29, 394-400.
- Muelbert, M. M., Bowen, W. D., Iverson, S. J., 2003. Weaning mass affects changes in body composition and food intake in harbour seal pups during the first month of independence. *Physiol. Biochem. Zool.* 76, 418-427.

- Newman, A. J., 1971. Cysts of branchial arch origin in the thymus of the beagle. *J. Small Anim. Pract.* 12, 681-685.
- Nielsen, J., 1995. Immunological and hematological parameters in captive harbor seals (*Phoca vitulina*). *Mar. Mamm. Sci.* 11, 314-323.
- Nieminen, P., Asikainen, J., Hyvarinen, H., 2001. Effects of seasonality and fasting on the plasma leptin and thyroxin levels of the racoon dog (*Nyctereutes procyonoides*) and the blue fox (*Alopex lagopus*). *J. Exp. Zool.* 289, 109-118.
- Nieminen, P., Saarela, S., Pyykonen, T., Asikainen, J., Mononen, J., Mustonen, A.-M., 2004. Endocrine response to fasting in the overwintering captive raccoon dog (*Nyctereutes procyonoides*). *J. Exp. Zool.* 301A, 919-929.
- Nilssen, K. T., Haug, T., Lindblom, C., 2001. Diet of weaned pups and seasonal variations in body condition of juvenile Barents Sea harp seals *Phoca groenlandica*. *Mar. Mamm. Sci.* 17, 926-936.
- Nunn, C. L., Gittleman, J. L., Antonvics, J., 2003. A comparative study of white blood cell counts and disease risk in carnivores. *Proc. R. Soc. Lond. B.* 270, 347-356.
- Nunn, C. L., Lindenfors, P., Pursall, E. R., Rolff, J., 2009. On sexual dimorphism in immune function. *Phil. Trans. R. Soc. B.* 364, 61-69.
- Oftedal, O. T., Boness, D. J., Tedman, R. A., 1987. The behavior, physiology, and anatomy of lactation in the pinnipedia. In: H. H. Genoways, (Ed.), *Current Mammalogy*. Plenum Publishing Corp, p. 175-245.
- Oki, C., Atkinson, S., 2004. Diurnal patterns of cortisol and thyroid hormones in the Harbor Seal (*Phoca vitulina*) during summer and winter seasons. *Gen. Com. Endocrinol.* 136, 289-297.
- Ortiz, R. M., Wade, C. E., Ortiz, C. L., 2000. Prolonged fasting increases the response of the Renin-Angiotensin-Aldosterone system, but not vasopressin levels, in postweaned northern elephant seal pups. *Gen. Comp. Endocrinol.* 119, 217-223.
- Ortiz, R. M., Noren, D. P., Litz, B., Ortiz, C. L., 2001. A new perspective on adiposity in a naturally obese mammal. *Am. J. Physiol. Endocrinol. Metab.* 281, E1347-E1351.
- Ortiz, R. M., Wade, C. E., Ortiz, C. L., Talamantes, F., 2003a. Acutely elevated vasopressin increases circulating concentrations of cortisol and aldosterone in fasting northern elephant seal (*Mirounga angustirostris*) pups. *J. Exp. Biol.* 206, 2795-2802.

- Ortiz, R. M., Houser, D. S., Wade, C. E., Ortiz, C. L., 2003b. Hormonal changes associated with the transition between nursing and natural fasting in northern elephant seals (*Mirounga angustirostris*). *Gen. Comp. Endocrinol.* 130, 78-83.
- Ortiz, R. M., Crocker, D. E., Houser, D. S., Webb, P. M., 2006. Angiotensin II and aldosterone increase with fasting in breeding adult male northern elephant seals (*Mirounga angustirostris*). *Physiol. Biochem. Zool.* 79, 1106-1112.
- Petrauskas, L., Atkinson, S., Gulland, F., Mellish, J.-A., Horning, M., 2008. Monitoring glucocorticoid response to rehabilitation and research procedures in California and Steller sea lions. *J. Exp. Zool.* 309A, 73-82.
- Pitcher, K., Calkins, D. G., Pendleton, G. W., 2000. Steller sea lion body condition indices. *Mar. Mamm. Sci.* 16, 427-436.
- Pitcher, K. W., 1990. Major decline in number of harbor seals, *Phoca vitulina richardsi*, on Tugidak Island, Gulf of Alaska. *Mar. Mamm. Sci.* 6, 121-134.
- Ponganis, P. J., Kooyman, G. L., Sartoris, D., Jobsis, P., 1997. Pinniped splenic volumes. *Am. J. Physiol. Regul. Integr. Comp. Physiol.* 82, 1989-1994.
- Rea, L. D., 2002. Indices of condition in Steller sea lions (*Eumetopias jubatus*). In: D. DeMaster, S. Atkinson, (Eds.), *Steller sea lion decline: is it food II*. University of Alaska Sea Grant, Fairbanks, pp. 29-33.
- Rea, L. D., Castellini, M. A., Fadely, B., Loughlin, T. R., 1998. Health status of young Alaska Steller sea lion pups (*Eumetopias jubatus*) as indicated by blood chemistry and hematology. *Comp. Biochem. Physiol. A.* 120, 617-623.
- Rea, L. D., Rosen, D. A. S., Trites, A. W., 2007. Utilization of stored energy reserves during fasting varies by age and season in Steller sea lions. *Can. J. Zool.* 85, 190-200.
- Rea, L. D., Berman-Kowalewski, M., Rosen, D. A. S., Trites, A. W., 2009. Seasonal differences in biochemical adaptation to fasting in juvenile and subadult Steller sea lions (*Eumetopias jubatus*). *Physiol. Biochem. Zool.* 82, 236-247.
- Renouf, D., Gales, R., Noseworthy, E., 1993. Seasonal variation in energy intake and condition of harp seals: Is there a harp seal morph? Problems for bioenergetic modelling. *J. Zool. Lond.* 230, 513-528.
- Ridgway, S. H., 1972. *Mammals in the sea: biology and medicine*. Charles C. Thomas, Springfield, IL.

- Riviere, J. E., Englehardt, F. R., Solomon, J., 1977. The relationship of thyroxine and cortisol to the moult of the harbor seal *Phoca vitulina*. *Gen. Comp. Endocrinol.* 31, 398-401.
- Romano, T. A., Felten, S. Y., Olschowka, J. A., Felten, D. L., 1993. A microscopic investigation of the lymphoid organs of the beluga, *Delphinapterus leucas*. *J. Morphol.* 215, 261-287.
- Romano, T. A., Keogh, M. J., Kelly, C., Feng, P., Berk, L., Schlundt, C. E., Carder, D. A., Finneran, J. J., 2004. Anthropogenic sound and marine mammal health: measures of the nervous and immune systems before and after intense sound exposure. *Can. J. Fish. Aquat. Sci.* 61, 1124-1134.
- Romero, L. M., Meister, C. J., Cyr, N. E., Kenagy, G. J., Wingfield, J. C., 2008. Seasonal glucocorticoid responses to capture in wild free-living mammals. *Am. J. Physiol. Regul. Integr. Comp. Physiol.* 294, R614-R622.
- Rosen, D. A. S., Renouf, D., 1997. Seasonal changes in blubber distribution in Atlantic harbor seals: indication of thermodynamic considerations. *Mar. Mamm. Sci.* 13, 229-240.
- Rosen, D. A. S., Renouf, D., 1998. Correlates of seasonal changes in metabolism in Atlantic harbour seals (*Phoca vitulina concolor*). *Can. J. Zool.* 76, 1520-1528.
- Rosen, D. A. S., Hastie, G. D., Trites, A. W., 2004. Searching for stress: hematologic indicators of nutritional inadequacies in Steller sea lions. *Proc. Comp. Nutr. Soc.* 5, 145-149.
- Rosen, D. A. S., Kumagai, S., 2008. Hormone changes indicate that winter is a critical period for food shortages in Steller sea lions. *J. Comp. Physiol. B.* 178, 573-583.
- Ross, P. S., Pohajdak, B., Bowen, W. D., Addison, R. F., 1993. Immune function in free-ranging harbor seal (*Phoca vitulina*) mothers and their pups during lactation. *J. Wildl. Dis.* 29, 21-29.
- Ross, P. S., Swart, R. L. D., Visser, I. K. G., Vedder, L. J., Murk, W., Bowen, W. D., Osterhaus, A. D. M. E., 1994. Relative immunocompetence of the newborn harbour seal, *Phoca vitulina*. *Vet. Immunol. Immunopathol.* 42, 331-348.
- Rousseau, K., Atcha, Z., Loudon, A. S. I., 2003. Leptin and seasonal mammals. *J. Neuroendocrinol.* 15, 409-414.

- Ryg, M., Smith, T. G., Oritsland, N. A., 1990. Seasonal changes in body mass and body composition of ringed seals (*Phoca hispida*) on Svalbard. *Can. J. Zool.* 68, 470-475.
- Sangalang, G. B., Freeman, H. C., 1976. Steroids in the plasma of the gray seal, *Halichoerus grypus*. *Gen. Comp. Endocrinol.* 29, 419-422.
- Scheffer, V. B., 1960. Weights of organs and glands in the northern fur seal. *Mammalia*. 24, 476-481.
- Schulz, T. M., Bowen, W. D., 2004. Pinniped lactation strategies: Evaluation of data on maternal and offspring life history traits. *Mar. Mamm. Sci.* 20, 86-114.
- Schumacher, U., Welsch, U., 1987. Histological, histochemical, and fine structural observations on the spleen of seals. *Am. J. Anat.* 179, 356-368.
- Schumacher, U., Horny, H.-P., Heidemann, G., Schultz, W., Welsch, U., 1990. Histopathological findings in harbour seals (*Phoca vitulina*) found dead on the German North Sea coast. *J. Comp. Path.* 102, 299-309.
- Sease, J. L., Taylor, W. P., Loughlin, T. R., Pitcher, K. W. 2001. Aerial and land-based surveys of Steller sea lions (*Eumetopias jubatus*) in Alaska, June and July 1999 and 2000. U.S. Dep. Commer., NOAA Tech. Memo. NMFS-AFSC-122, p 52.
- Simpson, J. G., Gardner, M. B., 1972. Comparative microscopic anatomy of selected marine mammals. In: S. H. Ridgway, (Ed.), *Mammals of the Sea: Biology and Medicine*, Thomas. pp. 298-418.
- Slijper, E. J., 1958. Organ weights and symmetry problems in porpoises and seals. *Arch. Neerl. Sci. Nat.* 1, 97-113.
- Small, R. J., Pendleton, G. W., Pitcher, K. W., 2003. Trends in abundance of Alaska harbor seals, 1983-2001. *Mar. Mamm. Sci.* 19, 344-362.
- Smith, R. J., Lavigne, D. M., 1994. Subspecific status of the freshwater harbor seal (*Phoca vitulina mellonae*): a re-assessment. *Mar. Mamm. Sci.* 10, 105-110.
- Smodlaka, H., Henry, R. W., Reed, R. B., 2009. Macroscopic anatomy of the great vessels and structures associated with the heart of the ringed seal (*Pusa hispida*). *Anat. Histol. Embryol.* 38, 161-168.

- Springer, A. M., Estes, J. A., Van Vliet, G. B., Williams, T. M., Doak, D. F., Danner, E. M., Forney, K. A., Pfister, B., 2003. Sequential megafaunal collapse in the North Pacific Ocean: an ongoing legacy of industrial whaling? *Proc. Natl. Acad. Sci. U.S.A.* 100, 12223-12228.
- St. Aubin, D. J., Geraci, J. R., 1986. Adrenocortical function in pinniped hyponatremia. *Mar. Mamm. Sci.* 2, 243-250.
- St. Aubin, D. J., Ridgway, S. H., Wells, R. S., Rhinehart, H., 1996. Dolphin thyroid and adrenal hormones: circulating levels in wild and semidomesticated *Tursiops truncatus*, and influence of sex, age, and season. *Mar. Mamm. Sci.* 12, 1-13.
- Stewardson, C. L., Hemsley, S., Meyer, M. A., Canfield, P. J., Maindonald, J. H., 1999. Gross and microscopic visceral anatomy of the male Cape fur seal, *Arctocephalus pusillus pusillus* (Pinnipedia: Otariidae), with reference to organ size and growth. *J. Anat.* 195, 235-255.
- Stratakis, C. A., Chrousos, G. P., 1995. Neuroendocrinology and pathophysiology of the stress system. *Ann. N. Y. Acad. Sci.* 771, 1-18.
- Sucheston, M. E., Cannon, M. S., 1980. Cortex of the suprarenal (adrenal) gland of *Phoca vitulina richardi*. *Ohio J. Sci.* 80, 140-144.
- Suster, S., Rosai, J., 1990. Histology of the normal thymus. *Am. J. Surg. Pathol.* 14, 284-303.
- Swart, R. L. D., Kluten, R. M. G., Huizing, C. J., Vedder, L. J., Reijnders, P. J. H., Visser, I. K. G., Uytdehaag, F. G. C. M., Osterhaus, A. D. M. E., 1993. Mitogen and antigen induced B and T cell responses of peripheral blood mononuclear cells from the harbour seal (*Phoca vitulina*). *Vet. Immunol. Immunopathol.* 37, 217-230.
- Temte, J. L., Bigg, M. A., Wiig, O., 1991. Clines revisited: the timing of pupping in the harbour seal (*Phoca vitulina*). *J. Zool. Lond.* 224, 617-632.
- Thompson, P. M., Tollit, D. J., Corpe, H. M., Reid, R. J., Ross, H. M., 1997. Changes in haematological parameters in relation to prey switching in a wild population of harbour seals. *Funct. Ecol.* 11, 743-750.
- Thomson, C. A., Geraci, J. R., 1986. Cortisol, aldosterone, and leucocytes in the stress response of bottlenose dolphins, *Tursiops truncatus*. *Can. J. Fish. Aquat. Sci.* 43, 1010-1016.

- Trillmich, F., 1990. The behavioral ecology of maternal effort in fur seals and sea lions. *Behav.* 114, 3-20.
- Trites, A. W., Jonker, R. A. H., 2000. Morphometric measurements and body condition of healthy and starving Steller sea lion pups (*Eumetopias jubatus*). *Aquat. Mamm.* 26, 151-157.
- Trites, A. W., Donnelly, C. P., 2003. The decline of Steller seal lions *Eumetopias jubatus* in Alaska: a review of the nutritional stress hypothesis. *Mamm. Rev.* 33, 3-28.
- Trites, A. W., Deecke, V. B., Gregr, E. J., Ford, J. K. B., Olesiuk, P. F., 2007a. Killer whales, whaling, and sequential megafaunal collapse in the North Pacific: a comparative analysis of the dynamics of marine mammals in Alaska and British Columbia following commercial whaling. *Mar. Mamm. Sci.* 23, 751-765.
- Trites, A. W., Miller, A. J., Alexander, M. A., Bograd, S. J., Calder, J. A., Capotondi, A., Coyle, K. O., Lorenzo, E. D., Finney, B. P., Gregr, E. J., Grosch, C. E., Hare, S. R., Hunt Jr, G. L., Jahncke, J., Kachel, N. B., Kim, H.-J., Ladd, C., Mantua, N. J., Marzban, C., Maslowski, W., Mendelssohn, R., Neilson, D. J., Okkonen, S. R., Overland, J. E., Reedy-Maschner, K. L., Royer, T. C., Schwing, F. B., Wang, J. X. L., Winship, A. J., 2007b. Bottom-up forcing and the decline of Steller sea lions (*Eumetopias jubatus*) in Alaska: assessing the ocean climate hypothesis. *Fish. Oceanogr.* 16, 46-67.
- Trumble, S. J., Castellini, M. A., 2001. Blood chemistry and morphometric comparisons between harbor seal pups from Tugidak Island and within Prince William Sound, Alaska: using cluster analysis to assess health status. In. Harbor Seal Investigations, Alaska Department of Fish and Game Annual Report, NOAA # NA87FX0300, p. 324-344.
- Trumble, S. J., Castellini, M. A., 2002. Blood chemistry, hematology, and morphology of wild harbor seal pups in Alaska. *J. Wildl. Manag.* 66, 1197-1207.
- Trumble, S. J., Castellini, M. A., Mau, T. L., Castellini, J. M., 2006. Dietary and seasonal influences on blood chemistry and hematology in captive harbor seals. *Mar. Mamm. Sci.* 22, 104-123.
- Vander, A. J., Sherman, J. H., Luciano, D. S., 1986. Human physiology: the mechanisms of body function. McGraw-Hill, New York.
- Ver Hoef, J. M., Frost, K. J., 2003. A bayesian hierarchical model for monitoring harbor seal changes in Prince William Sound, Alaska. *Environ. Ecol. Stat.* 10, 201-219.
- Weber, K. T., 2003. A neuroendocrine-immune interface. *Herz.* 28, 692-701.

- Welsch, U., Schwerfirm, S., Skirnisson, K., Schumacher, U., 1997. Histological, histochemical, and fine structural observations on the lymph node of the common seal (*Phoca vitulina*) and the grey seal (*Halichoerus grypus*). *Anat. Rec.* 247, 225-242.
- Westlake, R. L., O'Corry-Crowe, G. M., 2002. Macrogeographic structure and patterns of genetic diversity in harbor seals (*Phoca vitulina*) from Alaska to Japan. *J. Mammal.* 83, 1111-1126.
- Westly, H. J., Kelley, K. W., 1984. Physiologic concentrations of cortisol suppress cell-mediated immune events in the domestic pig. *Proc. Soc. Exp. Biol. Med.* 177, 156-164.
- Williams, T. M., Rtishauser, M., Long, B., Fink, T., Gafney, J., Mostman-Liwanag, H., Casper, D., 2007. Seasonal variability in Otariid energetics: implications for the effects of predators on localized prey resources. *Physiol. Biochem. Zool.* 80, 433-443.
- Woldstad, S., Jenssen, B. M., 1999. Thyroid hormones in grey seal pups (*Halichoerus grypus*). *Comp. Biochem. Physiol. A.* 122, 157-162.
- York, A. E., 1994. The population dynamics of northern sea lions, 1975-1985. *Mar. Mamm. Sci.* 10, 38-51.
- Zapol, W. M., Liggins, G. C., Schneider, R. C., Qvist, J., Snider, M. T., Creasy, R. K., Hochachka, P. W., 1979. Regional blood flow during simulated diving in the conscious Weddell seal. *J. Appl. Physiol.* 47, 968-973.
- Zenteno-Savin, T., Castellini, M. A., Rea, L. D., Fadely, B. S., 1997. Plasma haptoglobin levels in threatened Alaska pinniped populations. *J. Wildl. Dis.* 33, 64-71.

Chapter 2:

Steller sea lion (*Eumetopias jubatus*) pups undergo a decrease in circulating white blood cells and the ability of T cells to proliferate during early post-natal development¹

2.1 Abstract

Postnatal changes in circulating immune components and peripheral blood mononuclear cells (PBMC) proliferation were assessed in Steller sea lion (SSL; *Eumetopias jubatus*) pups. Blood samples were collected for complete blood cell counts including total and differential white blood cell (WBC) counts from 46 pups ranging in age from 5 to 38 days old. Total WBC and neutrophil counts decreased in association with increased age of the pups. The ability of PBMC to proliferate was assessed by *in vitro* exposure to concanavalin A (ConA) or lipopolysaccharide (LPS) in 21 pups ranging in age from 7 to 32 days old. All SSL pups responded to *in vitro* stimulation with ConA and LPS 055:B5 indicating peripheral T and B cells are capable of responding to an antigenic challenge. ConA- induced T cell proliferation decreased with age while there was no change in spontaneous proliferation of PBMC or B cells exposed to LPS. The decreases in total WBC, neutrophil counts and T cell proliferation indicates that SSL undergo a period of postnatal development in cell-mediated immune function which is comparatively longer than phocid pups and consistent with other otariids.

2.2 Introduction

The western population of Steller sea lions has declined by more than 80% since the 1970's (Loughlin, 1998; Sease *et al.*, 2001) and is listed as endangered under the Endangered Species Act. While the mechanistic cause of the decline is a matter of continuing debate (Springer *et al.*, 2003; Fritz and Hinckley, 2005; DeMaster *et al.*, 2006; Trites *et al.*, 2007; Atkinson *et al.*, 2008), studies have suggested a reduction in juvenile survival (York, 1994; Holmes and York, 2003). As a result, many studies have focused

¹Keogh, M.J., J. M. Maniscalco, and S. Atkinson. 2010. Vet. Immunol. Immunopathol. 137:298-304.

on SSL pups and juveniles (Castellini *et al.*, 1993; Rea *et al.*, 1998; Brandon *et al.*, 2005; Myers *et al.*, 2010).

Hematology has been routinely used to assess the health of individuals and populations of SSL (Castellini *et al.*, 1993; Rea *et al.*, 1998). Hematocrit (Hct), hemoglobin (Hb), white blood cell (WBC) and platelet counts (Plt) have been used to assess nutritional status in pinnipeds (Thompson *et al.*, 1997; Rosen *et al.*, 2004; Trumble *et al.*, 2006). While red blood cell (RBC), mean corpuscular hemoglobin concentration (MCHC), mean corpuscular volume (MCV), Hct, and Hb decreased following birth, the same parameters increased to adult levels between 5 and 9 month in association with the development of diving ability and oxygen storage in SSL pups and juveniles (Richmond *et al.*, 2005).

In addition to hematology, PBMC proliferation assays are increasingly used to assess immuno-competence in pinnipeds (Swart *et al.*, 1993; Ross *et al.*, 1994; Swart *et al.*, 1994; Levin *et al.*, 2005; Mori *et al.*, 2006); however, few studies have assessed postnatal changes (Ross *et al.*, 1993; Ross *et al.*, 1994; Beckmen *et al.*, 1999; Kakuschke *et al.*, 2008). Ross *et al.*, (1993; 1994) concluded that harbor seal (HS; *Phoca vitulina*) pups have a functionally competent immune system at birth based upon functional changes during the 15 days following birth including: 1) a rapid increase in total IgG in serum, 2) *in vivo* antibody response to immunization, and 3) PBMC proliferation increases in response to *in vitro* mitogen stimulation. These conclusions were supported by the higher PBMC proliferation in HS pups compared to adults by Kakuschke *et al.*, (2008). The authors suggest these findings are due to unique life history traits in HS including long gestational period, large birth size, and short lactation period (Bowen *et al.*, 1992; Bowen *et al.*, 2001). Similarly, SSL have relatively long gestational periods and large pups when compared on a mass-specific basis of the female (Kovacs and Lavigne, 1992), although SSL have long periods of dependency with lactation lasting a year or more (Boness and Bowen, 1996). In contrast to the rapid increase in Ig in HS pups, the transfer of maternal antibodies, whether by placenta or colostrum, from adult northern fur seals (NFS; *Callorhinus ursinus*) to pups was low, despite adult Ig levels

similar to terrestrial mammals (Cavagnolo and Vedros, 1979). These findings were further supported following a *Klebsiella pneumonia* epizootic event in New Zealand sea lions (NZSL; *Phocarcos hookeri*). Castinel *et al.*, (2008) found that maternal antibodies for *K. pneumonia* were not passively transferred to pups and seroconversion occurring a minimum of 2 months after exposure. These findings suggest that unlike HS, the immune system of sea lions have a longer postnatal development period.

Capture and handling methods routinely used during blood collection in wild mammals can result in elevated cortisol concentrations (Engelhard *et al.*, 2002; Romero *et al.*, 2008), which may alter hematological parameters (Castellini *et al.*, 1996; Cattet *et al.*, 2003). Elevated cortisol concentrations have been shown to suppress PBMC proliferation in piglets (Westly and Kelley, 1984) while having no influence on lambs (Minton and Blecha, 1990) or calves (Manak, 1986). Given the inherent requirements of handling and the potential influence of cortisol on hematological components and *in vitro* PBMC proliferation, we quantified serum cortisol concentrations in an effort to assess the impact of sampling methods on these parameters.

The aim of the current study was to assess postnatal changes in circulating immune components and *in vitro* PBMC proliferation while accounting for potential effects of handling. Previous studies suggest that unlike HS, the immune system of sea lions may not be fully competent at birth (Ross *et al.*, 1993; Ross *et al.*, 1994). Therefore, we hypothesized that PBMC of SSL pups will proliferate when exposed to *in vitro* mitogen stimulation and that proliferation will be influenced by age as has been shown in other pinnipeds.

2.3 Materials and Methods

2.3.1 Study Site and Animals

Chiswell Island is home to a small SSL rookery located within the Alaska Maritime National Wildlife Refuge System, used by about 90 breeding animals producing up to 80 pups per year (Maniscalco *et al.*, 2006). The Chiswell Island rookery has been observed via remotely operated video monitoring equipment since 1998

(Maniscalco *et al.*, 2006). A total of 46 pups were measured, weighed, and blood sampled (Table 2.1). SSL pups were captured by corralling them in a natural rock formation on the rookery and later taking them to sampling stations individually. Body mass (BM) was

Table 2.1 Mean (\pm SD) for age, body mass (BM), standard length (SL), and axillary girth (AG) for SSL pups by sex and year.

		n	Age (day)	BM (kg)	SL (cm)	AG (cm)
June 30, 2005	Male	17	16.5 ± 5.2	31.5 ± 3.9	106.8 ± 4.0	74.1 ± 5.2
	Female	5	14.0 ± 5.6	28.0 ± 4.7	102.7 ± 4.9	71.4 ± 6.4
July 3, 2007	Male	10	23.3 ± 7.1	34.6 ± 4.5	110.4 ± 4.4	79.8 ± 4.4
	Female	5	14.1 ± 5.0	26.9 ± 6.0	101.8 ± 5.8	71.7 ± 6.8
July 1, 2008	Male	4	18.7 ± 10.4	32.3 ± 7.9	106.5 ± 8.7	75.3 ± 10.0
	Female	5	13.1 ± 3.9	25.6 ± 6.7	101.8 ± 2.8	72.3 ± 9.1
All	Male	31	18.9 ± 7.1	32.6 ± 4.7	107.9 ± 5.0	76.1 ± 6.8
	Female	15	13.8 ± 4.5	26.8 ± 4.4	102.1 ± 4.3	70.7 ± 5.7

measured to the nearest tenth of a kilogram using a hanging electronic scale (FWC series 7, FlexWeigh, Santa Rosa, CA). Standard length (SL) was measured as a straight line from tip-of-nose to tip-of-tail while the pup was lying on a straight board. Axillary girth (AG) was measured using a tape measure after the pup had exhaled. Pups ($n=43$) were permanently marked by hot iron branding (Merrick *et al.*, 1995) under isoflurane (USP; Halocarbon Industries, River Edge, NJ) anesthesia (Heath *et al.*, 1997) with the remaining three pups having flipper tags attached in the axillary area of both fore-flippers. Branding pups in conjunction with the remote video monitoring program allowed for the identification of mother-pup pairs and determination of pup ages (to within ± 4 hrs) by association with naturally marked females that were tracked from the time they gave birth. Pups ranged in age from 5 to 38 days old at the time of sampling.

Blood samples (< 16 ml) were collected using standard aseptic techniques from the caudal gluteal vein with an 18 gauge needle directly into Vacuette® blood collection tubes. Samples were collected in collaboration with other research and branding efforts. In 2005 and 2008 pups were anesthetized with isoflurane (ISO) prior to blood collection

and pups were physically restrained (PR) in 2007 during blood collection. EDTA-treated blood tubes were collected for CBC counts including total and differential WBC counts. Serum separator tubes were collected and serum was stored at -80 °C until cortisol was quantified. Sodium heparin-treated tubes were collected in 2007 and 2008 for isolation of PBMC from 21 of the pups (10 female, 11 male). Blood tubes were kept upright and chilled until further processing in the lab (< 12 hrs).

2.3.2 Complete Blood Cell Count

CBC counts were determined using the Heska[®] CBC-Diff Veterinary Hematological System (Heska[®] Corporation, Loveland, CO). Total WBC, Hct, RBC, Hb, and Plt were measured and used to calculate the MCV, MCH, and MCHC. Blood smears were made from whole EDTA blood and stained with Wright-Giemsa (Dip Quick Stain, Jorgensen Laboratories, Loveland, CO). WBC differentials, including neutrophils, lymphocytes, monocytes and eosinophils were counted manually and reported as percentages and absolute values of differentials (cells/ μ l).

2.3.3 Cortisol Radioimmunoassay

Cortisol was quantified using a solid-phase radioimmunoassay (RIA) (Siemens, Los Angeles, CA) previously validated for use with unextracted SSL serum (Mashburn and Atkinson, 2004). The RIA was performed per manufacturer instructions with the exception that all volumes were halved and run in duplicate. Inter-assay coefficients of variation for two assay controls were 5.6 and 10.0%. Intra-assay coefficients of variation were < 5.0 % and assay sensitivity was 2.0 ng/ml.

2.3.4 PBMC Isolation and Proliferation

Sodium heparin blood tubes were centrifuged at 200 x g for 10 min and the buffy coat was re-suspended in freezing media comprised of RPMI 1640, 20% FBS (Hyclone, Thermo Fisher Scientific Inc., Logan, UT), and 10% DMSO (Sigma Chemical Co., St. Louis, MO). Cryopreserved WBC were thawed and washed twice with PBS (without

$\text{Ca}^{2+}/\text{Mg}^{2+}$) and re-suspended in RPMI 1640 supplemented with 10% FBS, 2 mM L-glutamine, penicillin (100 U/ml), and streptomycin (100 $\mu\text{g}/\text{ml}$). PBMC were isolated by density gradient centrifugation (Histopaque 1077; MP Biomedical, Solon, OH) for 20 min at 720 x g. The PBMC containing band was isolated, washed twice, and re-suspended in RPMI 1640 supplemented with 10% FBS, 2 mM L-glutamine, penicillin (100 U/ml), and streptomycin (100 $\mu\text{g}/\text{ml}$), 1 mM sodium pyruvate, 100 μM nonessential amino acids, 25 mM hepes and 0.05 mM 2-mercaptoethanol (Sigma Chemical Co., St. Louis, MO). All cell culture media and supplements, except for FBS, DMSO and 2-mercaptoethanol, were purchased from Gibco, Invitrogen Corp., Carlsbad, CA. Viability was >90% and assessed using the exclusion dye trypan blue (Sigma Chemical Co., St. Louis, MO).

PBMC were plated (2.0×10^5 cells/well) in 96-well flat-bottom tissue-culture plates (Falcon, Becton Dickinson, Franklin Lakes, NJ) and their proliferation assessed following exposure to one of two mitogens (ConA or LPS 055:B5; Sigma Aldrich, St. Louis, MO). PBMC were cultured at 37°C with 5% CO_2 in triplicate for each animal in each of the following treatment groups; 1) unexposed control, 2) ConA suboptimal (0.1 $\mu\text{g}/\text{ml}$), 3) ConA optimal (1.0 $\mu\text{g}/\text{ml}$), 4) LPS suboptimal (1.0 $\mu\text{g}/\text{ml}$) and 5) LPS optimal (100 $\mu\text{g}/\text{ml}$). ConA is a plant-derived mitogen and preferentially stimulates T cells (Barta and Barta, 1993) while the LPS from *Escherichia coli* is a B cell activator (Wechsler-Reya and Monroe, 1996). The use of suboptimal and optimal mitogen concentrations have been used in previous pinniped studies (Levin *et al.*, 2005; Mori *et al.*, 2006). Mitogens were reconstituted in RPMI 1640 supplemented with 10% FBS, 2 mM L-glutamine, penicillin (100 U/ml), and streptomycin (100 $\mu\text{g}/\text{ml}$). These conditions are similar to those previously used for pinnipeds (Levin *et al.*, 2005; Mori *et al.*, 2006).

All PBMC were assessed in two assays with PBMC collected from a captive adult male SSL as an inter-assay control. PBMC were incubated for 48 h after which 20 μl of bromodioxuridine (BrdU) was added to a final concentration of 10 μM BrdU and incubated for an additional 18 h for a total incubation period of 66 h. Proliferation was assessed with the Cell Proliferation Biotrack ELISA System version 2 (GE Healthcare,

Piscataway, NJ) per manufacturer's instructions using Dynatech MRX Revelation microplate reader (Dynatech Laboratories, Inc, Chantilly, VA) at 450 nm with a reference wavelength of 650 nm. Data are presented as mean of the triplicates for optical density (OD) with standard deviation and as a stimulation index (SI; mean OD of cells exposed to mitogen/mean OD of cells in media only).

2.3.5 Statistical Analysis

Data were analyzed with Systat 10 (Systat Software, Inc, Point Richmond, CA). The best model for each parameter was selected using a stepwise general linear model with an iterative process of comparing the full mixed effects model, which included the categorical variables of sex and anesthesia, age as a continuous variable, and all interaction terms. The full model was compared to reduced models, which included only statistically significant grouping variables and interactions. Three of the 13 hematological parameters (MCV, MCH, MCHC) showed differences between PR and ISO; however, differences were less than 5% and therefore the hematological parameters were not separated for further analysis. To evaluate the effect of pup handling, serum cortisol concentrations were regressed against either the time elapsed from the initial arrival on the rookery to the time when the individual pup was removed from the corral and taken for blood collection or the immune components measured in the current study. Values were considered statistically significant if $p \leq 0.05$.

2.4 Results and Discussion

The ability to acquire blood samples from wild SSL pups of known age is a unique opportunity, which enabled us to assess postnatal changes in circulating immune components and PBMC proliferation. Based on BM and morphometrics all pups sampled were in good body condition for their respective age and sex. There was no effect of sex ($p=0.338$), anesthesia ($p=0.171$) or age ($p=0.854$) on circulating cortisol levels. Cortisol concentrations decreased when regressed against the elapsed time between arrival on the rookery and removal of pup from the holding corral for blood collection ($p=0.009$),

indicating that the pups were affected by the initial disturbance of the rookery. However, no significant effects of sex, anesthesia or serum cortisol concentrations ($p>0.082$) were detected in unstimulated or mitogen stimulated PBMC.

Proliferation of unstimulated PBMC and PBMC exposed to LPS and the suboptimal ConA did not change with age (Table 2.2). In piglets (Hoskinson *et al.*, 1990)

Table 2.2 Mean (\pm SD) and range of cortisol, hematology and PBMC proliferation from SSL pups. Age effect was significant at $p<0.05$.

Parameter	n	Means		Ranges		Age
Cortisol (ng/ml)	46	136.8	\pm 32.4	86.7	– 216.0	$p=0.854$
WBC ($10^3/\mu\text{l}$)	42	13.1	\pm 2.8	7.3	– 19.6	$p<0.001$
Neutrophil ($10^3/\mu\text{l}$)	42	8.60	\pm 2.54	4.36	– 16.97	$p=0.001$
(% of WBC)	42	64.9	\pm 9.2	45.0	– 84.0	$p=0.076$
Lymphocyte ($10^3/\mu\text{l}$)	42	2.55	\pm 1.18	1.12	– 5.63	$p=0.871$
(% of WBC)	42	19.7	\pm 8.3	8.0	– 42.0	$p=0.128$
Monocytes ($10^3/\mu\text{l}$)	42	1.54	\pm 0.87	0.0	– 3.8	$p=0.698$
(% of WBC)	42	11.83	\pm 6.35	0.0	– 24.0	$p=0.481$
Eosinophil ($10^3/\mu\text{l}$)	42	0.44	\pm 0.38	0.0	– 1.16	$p=0.635$
(% of WBC)	42	3.5	\pm 3.1	0.0	– 10.0	$p=0.467$
N:L ratio	42	4.06	\pm 2.16	1.07	– 9.75	$p=0.042$
Hct (%)	42	34.6	\pm 3.1	29.7	– 43.8	$p=0.142$
MCV (fl)	42	100.0	\pm 3.6	93.4	– 108.2	$p=0.594$
RBC ($10^6/\mu\text{l}$)	42	4.3	\pm 3.5	3.0	– 4.3	$p=0.175$
Hb (g/dl)	42	12.5	\pm 1.1	10.7	– 15.0	$p=0.229$
MCH (pg)	42	36.3	\pm 1.5	32.5	– 40.0	$p=0.416$
MCHC (g/dl)	42	36.3	\pm 1.8	34.0	– 39.4	$p=0.774$
PLT ($10^3/\mu\text{l}$)	42	367.9	\pm 120.1	93.0	– 639.0	$p=0.496$
Unstimulated (OD)	21	0.300	\pm 0.029	0.240	– 0.350	$p=0.942$
ConA						
0.1 $\mu\text{g/ml}$ (OD)	21	0.313	\pm 0.040	0.240	– 0.370	$p=0.137$
0.1 $\mu\text{g/ml}$ (SI)	21	1.05	\pm 0.11	0.890	– 1.280	$p=0.137$
1.0 $\mu\text{g/ml}$ (OD)	21	0.842	\pm 0.784	0.310	– 3.130	$p=0.039$
1.0 $\mu\text{g/ml}$ (SI)	21	6.10	\pm 2.49	1.330	– 10.440	$p=0.038$
LPS 055:B5						
1.0 $\mu\text{g/ml}$ (OD)	21	0.408	\pm 0.131	0.238	– 0.661	$p=0.728$
1.0 $\mu\text{g/ml}$ (SI)	21	1.360	\pm 0.410	0.900	– 2.10	$p=0.670$
100 $\mu\text{g/ml}$ (OD)	21	0.636	\pm 0.278	0.238	– 1.210	$p=0.309$
100 $\mu\text{g/ml}$ (SI)	21	2.080	\pm 0.850	0.920	– 3.740	$p=0.293$

and dogs (Toman *et al.*, 2002; Faldyna *et al.*, 2005) high spontaneous proliferation of PBMC were observed during the first week after birth and subsequently decreased within four to six weeks. The cause of the high spontaneous proliferation was unclear but may be due to PBMC being immature or in an activated state (Hoskinson *et al.*, 1990). In the present study, PBMC proliferated when exposed *in vitro* to ConA and LPS compared to unstimulated PBMC, suggesting that peripheral T and B cells are capable of responding to an antigenic challenge as early as 5 days following birth. Further, T cell proliferation decreased with age when exposed to the optimal ConA (Table 2.2, Figure 2.1). Similar




Figure 2.1 PBMC proliferation responses from 21 SSL pups exposed to 1.0 µg/ml ConA. Results presented as (A) absorbency (OD) and (B) stimulation index (SI). Significant ($p \leq 0.05$) trend lines are presented.

decreases in T cell proliferation during the first month of life have been observed in dogs followed by a gradual increase after 6 weeks, remaining below proliferation levels of adult dogs until at least 3 months of age (Gerber and Brown, 1974; Toman *et al.*, 2002). Similarly, an initial decrease in proliferation was observed in piglets (Hoskinson *et al.*, 1990), rabbits (Jeklova *et al.*, 2007), and NFS pups (Beckmen, 1999).

Beckmen (1999) reported a decrease in T cell proliferation exposed to ConA in NFS sampled as neonates (<7 days) and again at an estimated 29-51 days old. Moreover,

NFS pups born to older females (43%) developed adequate antibody titers when vaccinated with tetanus toxoid compared to only 5% of pups born to young females (Beckmen *et al.*, 2003). The low humoral immune response in pups of young females was associated with milk comparatively lower in fat (Beckmen *et al.*, 1999) and higher organochlorine (OC) levels resulting in the higher OG levels in the blood of NFS pups (Beckmen *et al.*, 1999; Beckmen *et al.*, 2003).


Development of antibodies against tetanus is dependent on T cells (Willcox, 1975). Decrease in T cell proliferation following birth observed in the current study and in NFS (Beckmen, 1999) suggests that age at time of vaccination is another factor contributing to the low humoral immune response observed by Beckmen *et al.*, (2003). Only 23% of the vaccinated NFS pups showed a 1.7-fold or greater increase in serum tetanus antibody, indicating a low humoral response in both the number of seroconverted individuals and in the low level of tetanus antibodies in pups considered to have achieved seroconversion (Beckmen *et al.*, 2003). These findings are consistent with observations in NZSL following a *K. pneumonia* epizootic event (Castinel *et al.*, 2008). In our study, SSL pups had a decrease in T cell proliferation comparable to changes observed in NFS (Beckmen, 1999) suggesting that SSL pups would have a similarly low or delayed humoral immune response as observed in other otariid pups (Beckmen *et al.*, 2003; Castinel *et al.*, 2008).


There was not a significant effect of sex or anesthesia on total or differential WBC counts, while monocyte counts ($p = 0.020$, Figure 2.2I) decreased in association with increasing cortisol concentrations. However, cortisol concentrations ($p > 0.059$) did not affect total WBC, neutrophil, lymphocyte, eosinophil counts or CBC count parameters including Hct, RBC, Hb, or PLT, suggesting handling of pups did not alter circulating immune components. Total WBC and neutrophil counts decreased while there was no change in lymphocyte, monocyte, or eosinophil counts with age (Table 2.2, Figure 2.2). The neutrophil to lymphocyte ratio also decreased with age. The decreases in total WBC and neutrophil to lymphocyte ratio were driven by changes in neutrophils. In HS pups, neutrophils also decreased but no change was observed in total WBC counts

due to a concurrent increase in lymphocytes and monocytes (Ross *et al.*, 1994). Hasselmeier *et al.*, (2008) also found higher levels of lymphocytes and monocytes in pups compared to older HS. Mean total WBC count in the current study was higher than reported values for SSL pups and juveniles (Mellish *et al.*, 2006). The higher WBC count in our study is likely due to pup age. Mellish *et al.* (2006) sampled older SSL pups (6 to 10 months) and juveniles (1 to 3 y) concluding that pups had significantly lower WBC counts than juveniles. Taken with the WBC counts in the current study, young SSL pups have a higher WBC followed by a postnatal decrease remaining below the levels observed in juveniles. Differential WBC counts in our study were within ranges reported for healthy adult California sea lions, *Zalophus californianus* (Roletto, 1993) and SSL pups (Bishop and Morado, 1995). Specifically, Bishop and Morado (1995) found similar percentage of neutrophil (67.7 ± 6.6), lymphocyte (15.1 ± 5.6), monocyte (11.5 ± 4.8) , and eosinophils (3.5 ± 2.8) though the mean total WBC was lower than in our study. Difference in total WBC count is likely explained by methodological differences between studies, total WBC counts were based upon estimated WBC density in the blood smear slides in the Bishop and Morado (1995) study.

This study adds to the limited knowledge of the postnatal changes in SSL pups. The decreases in total WBC, neutrophil counts and T cell proliferation indicates that SSL undergo a period of postnatal development in cell-mediated immune function, which is comparatively longer than in phocid pups and consistent with other otariids. The postnatal changes in SSL pups may be associated with differences in disease susceptibility between pups and adults. Future studies are needed to assess when pups or juveniles reach immuno-competence and if the development of immune function is associated with disease susceptibility contributing to the reduction in juvenile survival suggested in some SSL populations (York, 1994; Holmes and York, 2003). Given the unique life history traits of pinnipeds, these findings are relevant for comparative immunology and have implications for the interpretation of hematological parameters such as total and differential WBC counts in studies conducted on free-ranging SSL pups.

Figure 2.2 (A) Total WBC counts ($10^3/\mu\text{l}$) from 42 pups during the 38 days following birth and differential counts for (B) neutrophils ($10^3/\mu\text{l}$); (C) lymphocytes ($10^3/\mu\text{l}$); (D) monocytes ($10^3/\mu\text{l}$); and (E) eosinophils ($10^3/\mu\text{l}$). (F) Total WBC counts ($10^3/\mu\text{l}$) (G) neutrophils ($10^3/\mu\text{l}$); (H) lymphocytes ($10^3/\mu\text{l}$); (I) monocytes ($10^3/\mu\text{l}$); and (J) eosinophils ($10^3/\mu\text{l}$) are also presented against level of circulating cortisol (ng/ml). Significant ($p \leq 0.05$) trend lines are presented.





2.5 Acknowledgements

We would like to thank the veterinary, research, and husbandry staff at the Alaska SeaLife Center (ASLC) and the U.S. Fish and Wildlife Service Alaska Maritime National Wildlife Refuge for allowing access to Chiswell Island. This research was authorized under US Marine Mammal Protection Act Permit 881-1890-02 and under a U.S. Fish and Wildlife Service Special Use Permit for access to Chiswell Island. Protocols were reviewed and approved by the Institutional Animal Care and Use Committee at ASLC (07-001). This project was supported by the ASLC's Steller Sea Lion Research Program with funds from National Marine Fisheries Service and by the Pollock Conservation Cooperative Research Center, School of Fisheries and Ocean Sciences, University of Alaska Fairbanks, Alaska. However, the findings and conclusions presented by the authors are their own and do not necessarily reflect the views or positions of the Center or the University of Alaska. Drs. M. Castellini, T. Hollmen, R.M. Ortiz, and J. Runstadler and two anonymous reviewers graciously provided comments on this manuscript.

2.6 Literature Cited

- Atkinson, S., DeMaster, D. P., Calkins, D. G., 2008. Anthropogenic causes of the western Steller sea lion *Eumetopias jubatus* population decline and their threat to recovery. *Mamm. Rev.* 38, 1-18.
- Barta, O., Barta, V., 1993. Lymphocyte transformation test. In: Barta, O. (Ed.), *Veterinary Clinical Immunology Laboratory. Monographs in Animal Immunology*, 2. Bar-Lab, Inc Blacksburg, p. B4-1-B4-17.
- Beckmen, K. B., Ylitalo, G. M., Towell, R. G., Krahn, M. M., O'Hara, T. M., Blake, J. E., 1999. Factors affecting organochlorine contaminant concentrations in milk and blood of northern fur seal (*Callorhinus ursinus*) dams and pups from St. George Island, Alaska. *Sci. Total Environ.* 231, 183-200.
- Beckmen, K. B., Blake, J. E., Ylitalo, G. M., Stott, J. L., O'Hara, T. M., 2003. Organochlorine contaminant exposure and associations with hematological and humoral immune functional assays with dam age as a factor in free-ranging northern fur seal pups (*Callorhinus ursinus*). *Mar. Poll. Bull.* 46, 594-606.

- Bishop, D. H., Morado, J. F., 1995. Results on blood cell morphology and differential blood cell counts from seventeen Steller sea lion *Eumetopias jubatus* pups. Dis. Aquat. Org. 23, 1-6.
- Boness, D. J., Bowen, W. D., 1996. The evolution of maternal care in pinnipeds. BioScience. 46, 645-654.
- Bowen, W. D., Oftedal, O. T., Boness, D. J., 1992. Mass and energy transfer during lactation in a small Phoci, the harbor seal (*Phoca vitulina*). Physiol. Zool. 65, 844-866.
- Bowen, W. D., Ellis, S. L., Iverson, S. J., Boness, D. J., 2001. Maternal effects on offspring growth rate and weaning mass in harbour seals. Can. J. Zool. 79, 1088-1101.
- Brandon, E. A. A., Calkins, D. G., Davis, R. W., 2005. Neonatal growth of Steller sea lion (*Eumetopias jubatus*) pups in Alaska. Fish. Bull. 103, 246-257.
- Castellini, J. M., Melselman, H. J., Castellini, M. A., 1996. Understanding and interpreting hematocrit measurements in pinnipeds. Mar. Mamm. Sci. 12, 251-264.
- Castellini, M. A., Davis, R. W., Loughlin, T. R., Williams, T. M., 1993. Blood chemistries and body condition of Steller sea lion pups at Marmot Island, Alaska. Mar. Mamm. Sci. 9, 202-208.
- Castinel, A., Kittelberger, R., Pomroy, W. E., Dulgnan, P. J., Chilvers, B. L., Wilkinson, I. S., 2008. Humoral immune response to *Klebsiella spp.* in New Zealand sea lions (*Phocarctos hookeri*) and the passive transfer of immunity to pups. J. Wildl. Dis. 44, 8-15.
- Cattet, M. R. L., Christison, K., Caulkett, N. A., Stenhouse, G. B., 2003. Physiologic responses of grizzly bears to different methods of capture. J. Wildl. Dis. 39, 649-654.
- Cavagnolo, R. Z., Vedros, N. A., 1979. Serum and colostrum immunoglobulin levels in the northern fur seal *Callorhinus ursinus*. Dev. Comp. Immunol. 3, 139-146.
- DeMaster, D., P., Trites, A. W., Clapham, P., Mizroch, S. A., Wade, P., Small, R. J., Hoef, J. V., 2006. The sequential megafaunal collapse hypothesis: testing with existing data. Prog. Oceanogr. 68, 329-342.

- Engelhard, G. H., Brasseur, S. M. J. M., Hall, A. J., Burton, H. R., Reijnders, P. J. H., 2002. Adrenocortical responsiveness in southern elephant seal mothers and pups during lactation and the effect of scientific handling. *J. Comp. Physiol. B.* 172, 315-328.
- Faldyna, M., Sinkora, J., Knotigova, P., Leva, L., Toman, M., 2005. Lymphatic organ development in dogs: major lymphocyte subsets and activity. *Vet. Immunol. Immunopathol.* 104, 239-247.
- Fritz, L. W., Hinckley, S., 2005. A critical review of the regime shift-"junk food"-nutritional stress hypothesis for the decline of the western stock of Steller sea lion. *Mar. Mamm. Sci.* 21, 476-518.
- Gerber, J. D., Brown, A. L., 1974. Effect of development and aging on the response of canine lymphocytes to phytohemagglutinin. *Infect. Immun.* 10, 695-699.
- Hasselmeier, I., Fonfara, S., Driver, J., Siebert, U., 2008. Differential hematology profiles of free-ranging, rehabilitated, and captive harbor seals (*Phoca vitulina*) of the German North Sea. *Aquat. Mamm.* 34, 149-156.
- Heath, R. B., DeLong, R., Jameson, V., Bradley, D., Spraker, T., 1997. Isoflurane anesthesia in free ranging sea lion pups. *J. Wildl. Dis.* 33, 206-210.
- Holmes, E. E., York, A. E., 2003. Using age structure to detect impacts on threatened populations: a case study with Steller sea lions. *Conserv. Biol.* 17, 1794-1806.
- Hoskinson, C. D., Chew, B. P., Wong, T. S., 1990. Age-related changes in mitogen-induced lymphocyte proliferation and polymorphonuclear neutrophil function in the piglet. *J. Anim. Sci.* 68, 2471-2478.
- Jeklova, E., Leva, L., Kudlackova, H., Faldyna, M., 2007. Functional development of immune response in rabbits. *Vet. Immunol. Immunopathol.* 118, 221-228.
- Kakuschke, A., Valentine-Thon, E., Fonfara, S., Griesel, S., Rosenberger, T., Siebert, U., Prange, A., 2008. Metal-induced impairment of the cellular immunity of newborn harbor seals (*Phoca vitulina*). *Arch. Environ. Contam. Toxicol.* 55, 129-136.
- Kovacs, K. M., Lavigne, D. M., 1992. Maternal investment in otariid seals and walruses. *Can. J. Zool.* 70, 1953-1964.
- Levin, M., De Guise, S. D., Ross, P. S., 2005. Association between lymphocyte proliferation and polychlorinated biphenyls in free-ranging harbor seal (*Phoca vitulina*) pups from British Columbia, Canada. *Environ. Toxicol. Chem.* 24, 1247-1252.

- Loughlin, T. R., 1998. The Steller sea lion: a declining species. *Biosph. Conserv.* 1, 91-98.
- Manak, R. C., 1986. Neonatal calf serum immunoregulation of lymphocyte blastogenic responses. *J. Anim. Sci.* 62, 759-771.
- Maniscalco, J. M., Parker, P., Atkinson, S., 2006. Interseasonal and interannual measures of maternal care among individual Steller sea lions (*Eumetopias jubatus*). *J. Mammal.* 87, 304-311.
- Mashburn, K. L., Atkinson, S., 2004. Evaluation of adrenal function in serum and feces of Steller sea lions (*Eumetopias jubatus*): influences of molt, gender, sample storage, and age on glucocorticoid metabolism. *Gen. Comp. Endocrinol.* 136, 371-381.
- Mellish, J.-A. E., Calkins, D. G., Christen, D. R., Horning, M., Rea, L. D., Atkinson, S. K., 2006. Temporary captivity as a research tool: comprehensive study of wild pinnipeds under controlled conditions. *Aquat. Mamm.* 32, 58-65.
- Merrick, R. L., Brown, D. C., Calkins, D. G., Loughlin, T. R., 1995. A comparison of Steller sea lion, *Eumetopias jubatus*, pup masses between rookeries with increasing and decreasing populations. *Fish. Bull.* 93, 753-758.
- Minton, J. E., Blecha, F., 1990. Effect of acute stressors on endocrinological and immunological functions in lambs. *J. Anim. Sci.* 68, 3145-3151.
- Mori, C., Morsey, B., Levin, M., Nambiar, P. R., De Guise, S. D., 2006. Immunomodulatory effects of *in vitro* exposure to organochlorines on T-cell proliferation in marine mammals and mice. *J. Toxicol. Environ. Health A.* 69, 283-302.
- Myers, M. J., Litz, B., Atkinson, S., 2010. The effects of age, sex, season and geographic region on circulating serum cortisol concentrations in threatened and endangered Steller sea lions (*Eumetopias jubatus*). *Gen. Comp. Endocrinol.* 165, 72-77.
- Rea, L. D., Castellini, M. A., Fadely, B., Loughlin, T. R., 1998. Health status of young Alaska Steller sea lion pups (*Eumetopias jubatus*) as indicated by blood chemistry and hematology. *Comp. Biochem. Physiol. A.* 120, 617-623.
- Richmond, J. P., Burns, J. M., Rea, L. D., Mashburn, K. L., 2005. Postnatal ontogeny of erythropoietin and hematology in free-ranging Steller sea lions (*Eumetopias jubatus*). *Gen. Comp. Endocrinol.* 141, 240-247.

- Roletto, J., 1993. Hematology and serum chemistry values for clinically healthy and sick pinnipeds. *J. Zoo Wildl. Med.* 24, 145-157.
- Romero, L. M., Meister, C. J., Cyr, N. E., Kenagy, G. J., Wingfield, J. C., 2008. Seasonal glucocorticoid responses to capture in wild free-living mammals. *Am. J. Physiol. Reg. Integr. Comp. Physiol.* 294, R614-R622.
- Rosen, D. A. S., Hastie, G. D., Trites, A. W., 2004. Searching for stress: hematologic indicators of nutritional inadequancies in Steller sea lions. *Proc. Comp. Nutr. Soc.* 5, 145-149.
- Ross, P. S., Pohajdak, B., Bowen, W. D., Addison, R. F., 1993. Immune function in free-ranging harbor seal (*Phoca vitulina*) mothers and their pups during lactation. *J. Wildl. Dis.* 29, 21-29.
- Ross, P. S., Swart, R. L. D., Visser, I. K. G., Vedder, L. J., Murk, W., Bowen, W. D., Osterhaus, A. D. M. E., 1994. Relative immunocompetence of the newborn harbour seal, *Phoca vitulina*. *Vet. Immunol. Immunopathol.* 42, 331-348.
- Sease, J. L., Taylor, W. P., Loughlin, T. R., Pitcher, K. W., 2001. Aerial and land-based surveys of Steller sea lions (*Eumetopias jubatus*) in Alaska, June and July 1999 and 2000. U.S. Dep. Commer., NOAA Tech. Memo. NMFS-AFSC-122, p 52.
- Springer, A. M., Estes, J. A., Van Vliet, G. B., Williams, T. M., Doak, D. F., Danner, E. M., Forney, K. A., Pfister, B., 2003. Sequential megafaunal collapse in the North Pacific Ocean: an ongoing legacy of industrial whaling? *Proc. Natl. Acad. Sci. U.S.A.* 100, 12223-12228.
- Swart, R. L. D., Kluten, R. M. G., Huizing, C. J., Vedder, L. J., Reijnders, P. J. H., Visser, I. K. G., Uytdehaag, F. G. C. M., Osterhaus, A. D. M. E., 1993. Mitogen and antigen induced B and T cell responses of peripheral blood mononuclear cells from the harbour seal (*Phoca vitulina*). *Vet. Immunol. Immunopathol.* 37, 217-230.
- Swart, R. L. D., Ross, P. S., Vedder, L. J., Timmerman, H. H., Heisterkamp, S., Loveren, H. V., Vos, J. G., Reijnders, P. J. H., Osterhaus, A. D. M. E., 1994. Impairment of immune function in harbor seals (*Phoca vitulina*) feeding on fish from polluted waters. *Ambio* 23, 155-159.
- Thompson, P. M., Tollit, D. J., Corpe, H. M., Reid, R. J., Ross, H. M., 1997. Changes in haematological parameters in relation to prey switching in a wild population of harbour seals. *Func. Ecol.* 11, 743-750.

- Toman, M., Faldyna, M., Knotigova, P., Pokorova, D., Sinkora, J., 2002. Postnatal development of leukocyte subset composition and activity in dogs. *Vet. Immunol. Immunopathol.* 87, 321-326.
- Trites, A. W., Miller, A. J., Alexander, M. A., Bograd, S. J., Calder, J. A., Capotondi, A., Coyle, K. O., Lorenzo, E. D., Finney, B. P., Gregr, E. J., Grosch, C. E., Hare, S. R., Hunt Jr, G. L., Jahncke, J., Kachel, N. B., Kim, H.-J., Ladd, C., Mantua, N. J., Marzban, C., Maslowski, W., Mendelssohn, R., Neilson, D. J., Okkonen, S. R., Overland, J. E., Reedy-Maschner, K. L., Royer, T. C., Schwing, F. B., Wang, J. X. L., Winship, A. J., 2007. Bottom-up forcing and the decline of Steller sea lions (*Eumetopias jubatus*) in Alaska: assessing the ocean climate hypothesis. *Fish. Oceanogr.* 16, 46-67.
- Trumble, S. J., Castellini, M. A., Mau, T. L., Castellini, J. M., 2006. Dietary and seasonal influences on blood chemistry and hematology in captive harbor seals. *Mar. Mamm. Sci.* 22, 104-123.
- Wechsler-Reya, R. J., Monroe, J. G., 1996. Lipopolysaccharide prevents apoptosis and induces responsiveness to antigen receptor cross-linking in immature B cells. *Immunol.* 89, 356-362.
- Westly, H. J., Kelley, K. W., 1984. Physiologic concentrations of cortisol suppress cell-mediated immune events in the domestic pig. *Proc. Soc. Exp. Biol. Med.* 177, 156-164.
- Willcox, H. N. A., 1975. Thymus dependence of the antibody response to tetanus toxoid in mice. *Clin. Exp. Immunol.* 22, 341-347.
- York, A. E., 1994. The population dynamics of northern sea lions, 1975-1985. *Mar. Mamm. Sci.* 10, 38-51.

Chapter 3:

Body condition and endocrine profiles of healthy Steller sea lion (*Eumetopias jubatus*) pups during the early postnatal period¹

3.1 Abstract

Body condition indices have been useful in accessing the health in domestic livestock and free ranging populations of terrestrial mammals. Given the high energy demand and rapid growth during the postnatal period of Steller sea lion (SSL; *Eumetopias jubatus*) pups, body condition was expected to be related to circulating concentrations of a suite hormones (cortisol, aldosterone, thyroxine, triiodothyronine, leptin) associated with changes in body composition over time in developing pinnipeds. Blood samples were collected from free ranging SSL pups of known ages and sex. A body condition index (BCI) previously developed for SSL pups based on mass and length relationship was applied to 61 SSL pups ranging in age from 5 to 38 days old. However, no relationship between BCI and circulating concentration of hormones was detected. Circulating concentrations of cortisol, total and free thyroxine, and total triiodothyronine decreased when regressed against the elapsed time between researchers' arrival on the rookery and removal of pup from the holding corral for blood collection. While the overall variation attributed to the rookery disturbance was low ($r^2 < 0.293$), it may be of significance for future studies on free ranging pinnipeds. SSL pups sampled in the present study were larger on average than predicted by growth rates previously reported for SSL pups. These findings suggest that pups sampled in the present study are from a healthy population of SSL. This study adds to the limited knowledge of the postnatal changes in free ranging SSL pups by providing concentration ranges of several hormones from an apparently healthy population.

¹Keogh, M. J., S. Atkinson, J. M. Maniscalco. J. Comp. Physiol. B. *in submission*

3.2 Introduction

Body condition indices (BCI) have been applied in livestock to assess health and production measures (Caldeira et al. 2007) and to free ranging populations of terrestrial mammals (Cattet et al. 2002). Increasingly there has been an interest in quantifying body condition in pinnipeds as a measure of health and fitness (Arnould and Hindell 1999; Arnould and Warneke 2002; Castellini and Calkins 1993; Jonker and Trites 2000; Pitcher et al. 2000; Rea 2002; Trites and Jonker 2000). BCI for Steller sea lions (SSL; *Eumetopias jubatus*) have been derived from a combination of mass, length and blubber thickness (Pitcher et al. 2000; Rea 2002; Trites and Jonker 2000). BCI previously developed for SSL pups based on mass and length relationship were able to distinguish between dead pups with and without fat stores (Trites and Jonker 2000); however, how the BCI relate to circulating hormone concentrations has not been explored in SSL pups.

Changes in body condition or mass have been associated with changes in circulating cortisol, thyroid hormones and leptin in several species of terrestrial mammals (Barboza et al. 2004; Buff et al. 2002; Caldeira et al. 2007; Concannon et al. 2001) and pinnipeds (Jeanniard du Dot et al. 2009; Ortiz et al. 2003). Decreases in lipid stores and mass were associated with an increase in circulating cortisol concentration and decrease in total thyroxine and triiodothyronine concentrations in SSL (Jeanniard du Dot et al. 2009; Rosen and Kumagai 2008). These previous studies were based on longitudinal sampling during periods of experimental food restriction or fasting in juvenile and sub-adult SSL. Given the high energy demand and rapid growth (0.23–0.48 kg/day) during the postnatal period of SSL pups (Brandon et al. 2005), it is reasonable to expect an association between body condition and circulating concentrations of hormones associated with nutritional status and fat mass (leptin), lipid and water metabolism (cortisol and aldosterone), and growth and metabolism (thyroxine and triiodothyronine).

Currently two distinct population segments (DPS) of SSL are recognized within US waters: the eastern and western DPS. The western DPS is listed as endangered under the Endangered Species Act (U.S. Federal Register 62:30772–30773) having declined by more than 80% since the 1970's (Loughlin 1998; Sease et al. 2001). The causes of the

population decline and failure to recover have been the focus of continuing research and debate (Atkinson et al. 2008; DeMaster et al. 2006; Fritz and Hinckley 2005; Springer et al. 2003; Trites et al. 2007), predominately focusing on one or a combination of several factors including fisheries and interspecies competition for resources leading to a reduction in quantity or quality of food, human disturbance, predation, and exposure to contaminants. However, there remains no clear reason for the observed decline, which places an even greater emphasis on assessing the physiological factors (i.e. endocrine profiles) that impact body condition, and thus, health of this vulnerable species.

The present study sampled pups from a SSL rookery located on the northwestern shore of Chiswell Island in the eastern Gulf of Alaska within the western DPS. Prior to the western DPS decline, Chiswell Island supported a dramatically larger population of approximately 2,000 SSL (Mathisen and Lopp 1963); currently about 90 breeding animals use the rookery producing up to 80 pups annually (Maniscalco et al. 2006). Blood samples were collected from SSL pups of known ages, sex and body condition providing an opportunity to measure circulating hormones during the early postnatal period, and to determine if these hormones could provide predictable markers for evaluating the body condition of pups to better assess the health of the population of SSL. Therefore, the aims of the present study were to quantify circulating concentrations of cortisol, aldosterone, total and free thyroxine (T_4), total triiodothyronine (T_3), and leptin in SSL pups, and to determine if these hormones were associated with body condition. We hypothesized circulating concentrations of cortisol and aldosterone to decrease while leptin and thyroid hormones increase with increasing BCI. Further, capture and handling methods routinely used during blood collection in wild mammals can result in elevated circulating concentrations of cortisol (Engelhard et al. 2002; Romero 2002), aldosterone (Thomson and Geraci 1986), and thyroid hormones (Gartner et al. 1980; St. Aubin and Geraci 1988). Therefore, given the inherent requirements of handling necessary for sampling free ranging pinnipeds, we assessed the impact of the rookery disturbance on the hormones quantified in the present study.

3.3 Materials and Methods

3.3.1 Animals and Body Condition

A total of 61 randomly selected pups (male=37, female=24) were measured, weighed, and blood sampled on June 30, 2005; July 3, 2007 and July 1, 2008 (Table 1). SSL pups were captured after clearing the rookery of all adult SSL and corralling the pups in a natural rock formation on the rookery. Following capture, pups were individually taken to blood sampling stations. The time from corralling to removal for sampling ranged from 1 hr and 20 min to 8 hrs and 40 min. Body mass was measured to the nearest tenth of a kilogram using a hanging electronic scale (FWC series 7, FlexWeigh, Santa Rosa, CA), standard length was measured as a straight line from tip-of-nose to tip-of-tail while the pup was lying on a straight board with its ventral surface down, and axillary girth was measured using a tape measure after the pup exhaled. BCI were calculated by applying the stoutness-index [$\text{body mass} / (-63.88 + 0.8966 * \text{standard length})$] proposed by Trites and Jonker (2000).

Pups were marked by either branding (Merrick et al. 1995) or flipper-tagging for further identification to determine age. Marking pups in conjunction with the remote video monitoring program allowed for the identification of mother-pup pairs and determination of pup ages (to within ± 4 hrs) by association with naturally marked females that were tracked from the time they gave birth. Pups ranged in age from 5 to 38 days old at the time of blood collection. All SSL were continuously observed during daylight hours by the remote video camera monitoring system at the Alaska SeaLife Center (Seward, AK) throughout the entire breeding season (Maniscalco et al. 2006). Pups ranged in age from 5 to 38 days at the time of blood collection.

3.3.2 Blood Collection

Blood samples (< 18 ml) were collected using standard aseptic techniques from the caudal gluteal vein directly into Vacuette® serum separator blood collection tubes. In 2005 and 2008 pups were anesthetized under isoflurane (USP; Halocarbon Industries, River Edge, NJ) (Heath et al. 1997) prior to blood collection, and in 2007, pups were

physically restrained during blood collection. Serum separator tubes were kept upright and on ice until further processing in the laboratory (< 12 hrs). Serum was thereafter stored at -80°C until analysis.

3.3.3 Hormone Radioimmunoassays

Commercially available radioimmunoassay (RIA) kits were used to quantify hormone concentrations. All samples were run in duplicate per manufacturer instructions except where noted below, and with the exception that all volumes were halved. Solid-phase RIA kits (Siemens, Los Angeles, CA) were used for quantifying aldosterone, cortisol, and thyroid hormones. Cortisol, thyroid hormones (total T₃, total and free T₄), and leptin were previously validated in our laboratory (Mashburn and Atkinson 2004, 2008; Myers et al. 2006) and aldosterone was validated in the present study. Pools of male and female SSL serum were serially diluted to determine degree of parallelism to the standard curve. Recovery of added aldosterone (25–1200 pg/ml) for males ($y = -20.474x + 161.79$, $r^2 = 0.991$) and for females ($y = -20.803x + 164.39$; $r^2 = 0.997$) was determined. Leptin was assayed with a double-antibody multi-species leptin RIA kit (Linco Research, St. Charles, MO) previously validated with the following additional modifications to the manufacturer's protocol: the incubation period with primary antibody was doubled to 48 hr and buffer was not added to samples during incubation with the primary antibody (Mashburn and Atkinson 2008). Inter-assay and intra-assay coefficients of variation were <10.0% for each hormone. Assay sensitivities were as follows: cortisol, 2.0 ng/ml; aldosterone, 11.0 pg/ml; total T₃, 0.07 ng/ml; total T₄, 2.5 ng/ml; and free T₄, 0.1 pg/ml; leptin, 0.5 ng/ml human equivalent (HE).

3.3.4 Statistical Analysis

The best model for each hormone was selected using a stepwise general linear model with an iterative process of comparing the full mixed effects model, which included the categorical variables sex and anesthesia /physical restraint, with BCI or age as a continuous variable, and all interaction terms. The full model was compared to

reduced models, which included only variables and interactions with a $p \leq 0.100$. If anesthesia /physical restraint were identified as a significant factor, then pups under anesthesia or physical restraint were analyzed separately for effects of sex, age or BCI. Aldosterone concentrations were log transformed prior to statistical analysis to normalize the data. Mean values are reported \pm standard deviation (SD) and results were considered statistically significant if $p \leq 0.050$. To evaluate the effect of handling on pups, serum hormone concentrations measured in the present study were regressed against the time elapsed from the initial arrival on the rookery to the time when the individual pup was removed from the corral and taken for blood collection. Data were analyzed with Systat 10 (Systat Software, Inc, Point Richmond, CA).

3.4 Results

3.4.1 Age and Body Mass

BCI was not related to pup age ($p=0.216$) nor was there an effect of sex ($p=0.647$). Female pups were smaller than males in body mass ($p<0.001$), standard length ($p<0.001$), and axillary girth ($p=0.015$). SSL pups increased in body mass ($p<0.001$), standard length ($p<0.001$), and axillary girth ($p<0.001$) with age (Fig. 1) though female pups (14.7 ± 4.9 days) were on average younger than male pups (18.4 ± 7.2 day) ($p=0.025$).

3.4.2 Hormonal Profiles and Relationships

Male pups (368.1 ± 253.3 pg/ml) had higher aldosterone concentrations ($p=0.015$) (Fig. 2a) than females (357.0 ± 301.0 pg/ml) and there was a significant interaction between sex and BCI ($p=0.013$). The difference between male (323.5 ± 143.5 pg/ml) and female (276.5 ± 111.9 pg/ml) pups remained significant even when the outliers were removed ($p=0.011$). No significant effect of age or anesthesia /physical restraint was detected. Nor were aldosterone concentrations effected by the elapsed time between arrival on the rookery and blood collection. Circulating cortisol concentrations were higher in female (151.4 ± 45.1 ng/ml) than male (129.8 ± 34.0 ng/ml) pups ($p=0.037$)

(Fig. 2b) but were not affected by anesthesia /physical restraint, BCI or age. Cortisol concentrations decreased when regressed against the elapsed time between researchers arrival on the rookery and removal of pup from the holding corral for blood collection ($p < 0.001$) (Fig. 3a).

Leptin (ng/ml HE) serum levels were significantly higher ($p = 0.009$) in PR (2.0 ± 0.4 ng/ml HE) than pups under ISO (1.7 ± 0.4 ng/ml HE) during blood collection. When pups under anesthesia or physical restraint were analyzed separately, there was no effect of BCI, sex or age on circulating leptin concentrations.




Fig. 3.1. A total of 61 pups (male=37, female=24) were sampled during 2005, 2007, and 2008. Morphometrics for individual pups at age are presented in (a) body condition index, (b) axillary girth, (c) standard length, and (d) body mass. Female pups are represented by (●) and male pups are by (○).

Table 3.1 Mean (\pm SD) for age, body mass, standard length, and axillary girth for SSL pups by year and sex.

		n	Age (day)		n	body mass (kg)		n	standard length(cm)		n	axillary girth (cm)	
2005	Male	20	16.2	\pm 5.3	20	31.2	\pm 3.9	20	106.7	\pm 3.7	20	73.6	\pm 5.4
	Female	8	15.1	\pm 4.5	8	27.5	\pm 3.8	8	102.6	\pm 3.7	8	70.7	\pm 5.1
2007	Male	13	21.6	\pm 8.1	13	33.8	\pm 4.5	13	110.4	\pm 4.5	13	78.7	\pm 4.9
	Female	8	16.3	\pm 5.5	9	27.9	\pm 4.5	9	102.8	\pm 4.6	9	72.5	\pm 5.2
2008	Male	4	18.6	\pm 10.4	4	32.2	\pm 7.9	4	106.5	\pm 8.7	4	75.2	\pm 10.0
	Female	6	12.2	\pm 4.2	7	24.3	\pm 3.2	7	99.0	\pm 5.3	6	66.7	\pm 4.8
All	Male	37	18.3	\pm 7.2	37	32.2	\pm 4.6	37	108.0	\pm 4.9	37	75.6	\pm 6.1
	Female	22	14.7	\pm 4.9	24	26.7	\pm 4.1	24	101.6	\pm 4.7	23	70.3	\pm 5.4

* note: Age could not be determined from one female pup from 2007 and 2008 each.

There was no detectable effect of BCI, sex or ISO/PR on circulating total T_4 or free T_4 in SSL pups. Circulating total T_4 ($p=0.015$) and free T_4 ($p=0.031$) decreased with pup age (Fig. 3.4). Pups under anesthesia (0.6 ± 0.1 pg/ml) had higher ($p=0.001$) total T_3 compared to pups under physical restraint (0.5 ± 0.1 pg/ml) during blood collection. When compared separately there was no significant effect of sex, BCI or age in total T_3 . Thyroid hormones significantly decreased when regressed against the elapsed time from the initial arrival of researchers on the rookery to the time when the individual pup was removed from the corral and taken for blood collection (Fig. 3.3) including total T_4 ($p=0.008$), total T_3 ($p=0.012$), and free T_4 ($p<0.001$).




Fig. 3.2 (a) Aldosterone concentrations (pg/ml) and (b) cortisol concentrations (ng/ml) from female (●) and male (○) SSL pups. Dashed line separates the pups with greater than 2 SD of the mean for male and female pups for aldosterone concentration (pg/ml).

Table 3.2 Mean (\pm SD) for cortisol, aldosterone, total and free thyroxine (T₄), total triiodothyronine (T₃), and leptin concentrations for all 61 SSL pups and separated by sex (female=24, male=37).

	All pups			Females			Males		
	Mean \pm SD			Mean \pm SD			Mean \pm SD		
cortisol (ng/ml)*	138.3	\pm	39.8	151.4	\pm	45.1	129.8	\pm	33.9
aldosterone (pg/ml)*	364.2	\pm	270.6	357.0	\pm	300.9	368.8	\pm	253.3
total T ₄ (ng/ml)	19.5	\pm	5.3	20.0	\pm	5.3	19.2	\pm	5.3
total T ₃ (ng/ml)	0.6	\pm	0.1	0.6	\pm	0.2	0.5	\pm	0.1
free T ₄ (pg/ml)	6.1	\pm	2.1	6.1	\pm	1.6	6.1	\pm	2.3
total T ₄ :free T ₄	3.5	\pm	1.3	3.4	\pm	1.2	3.6	\pm	1.3
leptin (ng/ml HE)	1.9	\pm	0.4	1.9	\pm	0.4	1.7	\pm	0.4

*Statistically different between male and female pups at $p < 0.050$.




Fig 3.3 Circulating concentrations of (a) cortisol (ng/ml); (b) total T_4 (ng/ml); (c) free T_4 (pg/ml); (d) total T_3 (ng/ml) from female (●) and male (○) SSL pups regressed against the elapsed time from the initial arrival on the rookery to the time when individual pups were removed from the corral for blood collection. Female pups are represented by filled circles, males are by hollow circles. Significant ($p \leq 0.05$) trend lines are presented.




Fig 3.4. Circulating thyroid hormone concentrations from female (●) and male (○) SSL pups. (a, b) total T₄ (ng/ml); (c, d) free T₄ (pg/ml); (e, f) total T₃ (ng/ml) presented against the body condition index (BCI) or age. Female pups are represented by filled circles, males are by hollow circles. Only significant ($p \leq 0.05$) trend lines are presented.

3.5 Discussion

Changes in body condition and mass and associated changes in circulating cortisol, thyroid hormones and leptin have been well documented in several species of terrestrial mammals (Barboza et al. 2004; Buff et al. 2002; Caldeira et al. 2007; Concannon et al. 2001) and pinnipeds (Jeanniard du Dot et al. 2009; Ortiz et al. 2003). However, we found no relationship between BCI and any of the circulating concentrations of the hormones quantified in the present study on SSL. Further, body mass and a density index calculated as $[\text{body mass} / (\text{standard length} \times \text{axillary girth}^2)] \times 10^6$ (Castellini et al. 1993; Rea 2002) were similarly not related to circulating concentrations of any hormones assessed in the present study (data not shown).

In the present study, pups were in a period of rapid growth as evident by the increase in body mass, standard length and axillary girth with age. Males between 5 and 38 days of age were larger than females in body mass, standard length and axillary girth. These findings have been previously reported for SSL pups (Brandon et al. 2005). Brandon et al. (2005) found that during the first six weeks following birth, male and female SSL pups grew at the same rate in body mass, standard length and axillary girth and suggested that males were larger than females at birth by approximately 15%. Estimated mass at birth (female 18.7 kg, male 22.4 kg) did not vary between rookeries while growth rates in body mass ranged from 0.23 kg/day to 0.48 kg/day with higher rates found in the Aleutian Islands (Brandon et al. 2005). Pups from the Chiswell Island rookery were not included in that study and Chiswell Island is geographically closer to rookeries with the lower growth rates. Yet, if we apply the growth rates calculated by Brandon et al. (2005) to SSL pups of known ages in the present study, then SSL pups from Chiswell Island rookery are estimated to be 4.1 and 17.9% (females) and 3.0 and 17.2% (males) larger than predicted by the lowest and maximum estimated growth rates, respectively. Further, pups in the present study on average weighed slightly more than predicated for females (1.96%) and males (2.42%) based on the BCI proposed by Trites and Jonker (2000). These findings are likely an indication that pups sampled in the present study are from a healthy population of SSL. Further, there were no abandoned or

starving pups observed on Chiswell Island at the time of blood collection (J. Maniscalco, personal communication).

While we detected a significant difference in size between male and female pups, suggesting either a difference in sex-related physiology or maternal investment during gestation as Brandon et al. (2005) proposed, only cortisol and aldosterone showed differences between male and female pups. Aldosterone is integral to the regulation of water and sodium balance in mammals. Within phocids, aldosterone concentrations significantly increased during extended periods of fasting in pups (Englehardt and Ferguson 1980; Nordoy et al. 1993; Ortiz et al. 2000). Unlike phocids, SSL pups have short periods of fasting associated with the maternal foraging trips, which tend to increase in length as the pup ages (Higgins et al. 1988; Maniscalco et al. 2006; Trites and Porter 2002). However, the large variation in aldosterone concentrations observed in the present study was not associated with age or BCI of the SSL pup and the difference between male and female pups, while statistically significant, was small and may not be biologically relevant. Cortisol is well known as a stress hormone and plays a role in gluconeogenesis and promoting the mobilization of fatty acids from peripheral adipose tissues. Cortisol concentrations in the present study increased as a result of the initial disturbance of the rookery and continually decreased as the time elapsed between the initial disturbance and collection of blood. Cortisol concentrations at the time of initial capture in free ranging juvenile SSL (Mellish et al. 2007) were comparable to pups in the present and previous studies (Myers et al. 2010). Similar to the present study, Myers et al. (2010) reported cortisol concentrations in female pups (155.2 ± 2.3 ng/ml) higher than male pups (141.5 ± 2.1 ng/ml). Higher baseline GC or metabolite concentrations have also been reported in adult female terrestrial mammals (Franceschini et al. 2007) and females exposed to an acute stressor exhibited higher peak cortisol concentrations than males (Guimont and Wynne-Edwards 2006). While the large sample size in the present study may have allowed for the detection of significant differences between male and female pups, the biological relevance of the small difference in cortisol and aldosterone concentrations between male and female pups is unclear.

Thyroid hormones including total and free T_4 and total T_3 are integral to maintaining thermoregulation, metabolism and the growth and development of neonatal and juvenile mammals. All forms of thyroid hormones quantified decreased when regressed against elapsed time since arrival on rookery, although the overall variation attributed to the elapsed time was low ($r^2 < 0.293$). The decrease in circulating total and free T_4 and total T_3 could be the result of the initial rookery disturbance and associated increased activity of pups leading to elevated concentrations of thyroid hormones followed by a period of decreasing concentrations as circulating hormones return to basal concentrations. Alternatively, adrenocorticotrophic hormone and glucocorticoids have been shown to alter the secretion of thyroid-stimulating hormone from the pituitary (Dussault 1974; Gartner et al. 1980; Wilber and Utiger 1969) and the conversion of T_4 to T_3 in tissues (Heyma and Larkins 1982; Kaplan and Utiger 1978) leading to reduced concentrations of thyroid hormones. Further, the elapsed time is by necessity later in the day and given the role of thyroid hormones in regulation of metabolism and thermoregulation, the decrease in thyroid hormones may not solely be related to the rookery disturbance but rather the result of increasing temperatures as the day progressed. Given the experimental design of the present study we are unable to determine the direct cause of the observed decrease in thyroid hormone concentrations over the elapsed time between arrival on rookery and sampling.

Mean thyroid hormone concentrations (total and free T_4 , total T_3) were similar to those previously reported for SSL pups (Myers et al. 2006). However, while Myers et al. (2006) reported undetectable levels for some pups for all forms of thyroid hormones quantified, we did not have any SSL pups with undetectable levels for total and free T_4 or total T_3 but we were unable to quantify free T_3 (data not shown). Serum total T_3 was slightly lower in SSL pups but within the range reported in juvenile and sub-adult SSL (Jeanniard du Dot et al. 2009; Rosen and Kumagai 2008). SSL pups had total T_4 concentrations more than two times greater than baseline concentration during summer in juvenile and sub-adult SSL (Jeanniard du Dot et al. 2009; Rosen and Kumagai 2008) while free T_4 concentrations in SSL pups were half the circulating concentration found in

juvenile and sub-adult SSL (Jeanniard du Dot et al. 2009). Both total and free T_4 concentrations decreased with pup age, and while significant, the relationship was extremely weak ($r^2=0.098$) likely due to the short developmental period (5-38 days) covered in the present study. While male pups were larger than female pups at the time of sampling, no difference in circulating concentrations of any forms of thyroid hormones quantified was observed between male and female pups. These findings were not unexpected since Brandon et al. (2005) determined male and female pups have the same postnatal growth rate (body mass, standard length, and axillary girth) and concluded the difference in size between male and female pups was the result of differences in maternal investment during gestation.

Leptin is secreted by adipose tissue (Friedman and Halaas 1998; Kershaw and Flier 2004) and circulating concentrations have been correlated with fat stores and respond to changes in energy balance in terrestrial mammals (Friedman and Halaas 1998). We found higher leptin concentrations in pups physically restrained during blood collection compared to those under isoflurane. This finding was unexpected but driven by differences in fat stores or body mass as fluctuating leptin concentrations not associated with changes in fat or body mass have been previously reported in pinnipeds (Arnould et al. 2002; Mashburn and Atkinson 2008). In addition, the leptin concentrations in our study were near the lower detection limit of the kit (0.5 ng/ml) and similar to previous studies in phocids with similar methods (Hammond et al. 2005). Difficulty in quantifying leptin in SSL (Rosen and Kumagai 2008) and other pinnipeds (Arnould et al. 2002; Ortiz et al. 2003) has been previously reported with low concentrations or no correlation to body mass or lipid stores being reported (Arnould et al. 2002; Ortiz et al. 2001; Rosen and Kumagai 2008).

Circulating concentrations of cortisol, total and free T_4 , and total T_3 decreased when regressed against the elapsed time between researchers' arrival on the rookery and removal of pup from the holding corral for blood collection. While the overall variation attributed to the rookery disturbance was low it may be of significance for future studies on free ranging pinnipeds. While we did not find BCI to be related to circulating hormone

concentrations future studies should include pups with a wider range of body conditions including abandoned or starving pups and ideally would employ a longitudinal sampling scheme to address some of the large individual variation in hormone concentrations observed in the present study. This study adds to the limited knowledge of the postnatal changes in free ranging SSL pups by providing concentration ranges of several hormones from an apparently healthy population. Further, given the unique life history traits of pinnipeds, these findings are also relevant for comparative endocrinology.

3.6 Acknowledgements

We would like to thank the veterinary and research staff at the Alaska SeaLife Center (ASLC) and the University of Alaska School of Fisheries and Ocean Sciences Endocrine Laboratory. Drs. M. Castellini, T. Hollmen, R.M. Ortiz, J. Runstadler graciously provided comments on this manuscript. This project was supported by the Pollock Conservation Cooperative Research Center, School of Fisheries and Ocean Sciences, University of Alaska Fairbanks, Alaska and the ASLC's Steller Sea Lion Research Program with funds from National Marine Fisheries Service. However, the findings and conclusions presented by the authors are their own and do not necessarily reflect the views or positions of the Center or the University of Alaska. Additional funding was provided through a fellowship from the National Science Foundation GK-12 award to the University of Alaska Fairbanks "New GK-12 Program: The CASE (Changing Alaska Science Education) for Enhancing Understanding of Climate Change". This research was authorized under US Marine Mammal Protection Act Permit 881-1890-02 and under a U.S. Fish and Wildlife Service Special Use Permit for access to Chiswell Island. Protocols were reviewed and approved by the Institutional Animal Care and Use Committee at ASLC (07-001).

3.7 Literature Cited

- Arnould JPY, Hindell MA (1999) The composition of Australian fur seal (*Arctocephalus pusillus doriferus*) milk throughout lactation. *Physiol Biochem Zool* 72:605-612
- Arnould JPY, Morris MJ, Rawlins DR, Boyd IL (2002) Variation in plasma leptin levels in response to fasting in Antarctic fur seals (*Arctocephalus gazella*). *J Comp Physiol B* 172:27-34
- Arnould JPY, Warneke RM (2002) Growth and condition in Australian fur seals (*Arctocephalus pusillus doriferus*) (Carnivora:Pinnipedia). *Aust J Zool* 50:53-66
- Atkinson S, DeMaster DP, Calkins DG (2008) Anthropogenic causes of the western Steller sea lion *Eumetopias jubatus* population decline and their threat to recovery. *Mamm Rev* 38:1-18
- Barboza PS, Hartbauer DW, Hauer WE, Blake JE (2004) Polygynous mating impairs body condition and homeostasis in male reindeer (*Rangifer tarandus tarandus*). *J Comp Physiol B* 174:309-317
- Brandon EAA, Calkins DG, Davis RW (2005) Neonatal growth of Steller sea lion (*Eumetopias jubatus*) pups in Alaska. *Fish Bull* 103:246-257
- Buff PR, Dodds AC, Morrison CD, Whitley NC, McFadin EL, Daniel JA, Djiane J, Keisler DH (2002) Leptin in horses: tissue localization and relationship between peripheral concentrations of leptin and body condition. *J Anim Sci* 80:2942-2948
- Caldeira RM, Belo AT, Santos CC, Vazques MI, Portugal AV (2007) The effect of body condition score on blood metabolites and hormonal profiles in ewes. *Small Rumin Res* 68:233-241
- Castellini MA, Calkins DG (1993) Mass estimates using body morphology in Steller sea lions. *Mar Mamm Sci* 9:48-54
- Castellini MA, Davis RW, Loughlin TR, Williams TM (1993) Blood chemistries and body condition of Steller sea lion pups at Marmot Island, Alaska. *Mar Mamm Sci* 9:202-208
- Cattet MRL, Caulkett NA, Obbard ME, Stenhouse GB (2002) A body-condition index for ursids. *Can J Zool* 80:1156-1161
- Concannon P, Levac K, Rawson R, Tennant B, Bensadoun A (2001) Seasonal changes in serum leptin, food intake, and body weight in photoentrained woodchucks. *Am J Physiol Regul Integr Comp Physiol* 281:R951-R959

- DeMaster DP, Trites AW, Clapham P, Mizroch SA, Wade P, Small RJ, Hoef JV (2006) The sequential megafaunal collapse hypothesis: testing with existing data. *Prog Oceanogr* 68:329-342
- Dussault JH (1974) The effect of dexamethasone on TSH and prolactin secretion after TRH stimulation. *Can Med Assoc J* 111:1195-1197
- Engelhard GH, Brasseur SMJM, Hall AJ, Burton HR, Reijnders PJH (2002) Adrenocortical responsiveness in southern elephant seal mothers and pups during lactation and the effect of scientific handling. *J Comp Physiol B* 172:315-328
- Englehardt FR, Ferguson JM (1980) Adaptive hormone changes in harp seals, *Phoca groenlandica*, and gray seals, *Halichoerus grypus*, during the postnatal period. *Gen Comp Endocrinol* 40:434-445
- Franceschini C, Siutz C, Palme R, Millesi E (2007) Seasonal changes in cortisol and progesterone secretion in Common hamsters. *Gen Comp Endocrinol* 152:14-21
- Friedman JM, Halaas JL (1998) Leptin and the regulation of body weight in mammals. *Nat* 395:763-770
- Fritz LW, Hinckley S (2005) A critical review of the regime shift-"junk food"- nutritional stress hypothesis for the decline of the western stock of Steller sea lion. *Mar Mamm Sci* 21:476-518
- Gartner K, Buttner D, Dohler K, Friedel R, Lindena J, Trautschold I (1980) Stress response of rats to handling and experimental procedures. *Lab Anim* 14:267-274
- Guimont FS, Wynne-Edwards KE (2006) Individual variation in cortisol responses to acute "on-back" restraint in an outbred hamster. *Horm Behav* 50:252-260
- Hammond JA, Bennett KA, Walton MJ, Hall AJ (2005) Molecular cloning and expression of leptin in gray and harbor seal blubber, bone marrow, and lung and its potential role in marine mammal respiratory physiology. *Am J Physiol Regul Integr Comp Physiol* 289:R545-R553
- Heath RB, DeLong R, Jameson V, Bradley D, Spraker T (1997) Isoflurane anesthesia in free ranging sea lion pups. *J Wildl Dis* 33:206-210
- Heyma P, Larkins RG (1982) Glucocorticoids decrease in conversion of thyroxine into 3,5,3'-triiodothyronine by isolated rat renal tubules. *Clin Sci* 62:215-220

- Higgins LV, Costa DP, Huntley AC, Le Boeuf BJL (1988) Behavioral and physiological measurements of maternal investment in the Steller sea lion, *Eumetopias jubatus*. Mar Mamm Sci 4:44-58
- Jeanniard du Dot T, Rosen DAS, Richmond JP, Kitaysky AS, Zinn SA, Trites AW (2009) Changes in glucocorticoid, IGF-I and thyroid hormones as indicators of nutritional stress and subsequent refeeding in Steller sea lions (*Eumetopias jubatus*). Comp Biochem Physiol A 152:524-534
- Jonker RAH, Trites AW (2000) The reliability of skinfold-calipers for measuring blubber thickness of Steller sea lion pups (*Eumetopias jubatus*). Mar Mamm Sci 16:757-766
- Kaplan MM, Utiger RD (1978) Iodothyronine metabolism in rat liver homogenates. J Clin Investig 61:459-471
- Kershaw EE, Flier JS (2004) Adipose tissue as an endocrine organ. J Clin Endocrinol Metab 89:2548-2556
- Loughlin TR (1998) The Steller sea lion: a declining species. Biosph Conserv 1:91-98
- Maniscalco JM, Parker P, Atkinson S (2006) Interseasonal and interannual measures of maternal care among individual Steller sea lions (*Eumetopias jubatus*). J Mammal 87:304-311
- Mashburn KL, Atkinson S (2004) Evaluation of adrenal function in serum and feces of Steller sea lions (*Eumetopias jubatus*): influences of molt, gender, sample storage, and age on glucocorticoid metabolism. Gen Comp Endocrinol 136:371-381
- Mashburn KL, Atkinson S (2008) Variability in leptin and adrenal response in juvenile Steller sea lions (*Eumetopias jubatus*) to adrenocorticotrophic hormone (ACTH) in different seasons. Gen Comp Endocrinol 155:352-258
- Mathisen OA, Lopp RJ (1963) Photographic census of the Steller sea lion herds in Alaska, 1956-58. United States Fish and Wildlife Service, Special Scientific Report of Fisheries. 424: 1-20
- Mellish J-AE, Hennessy K, Thomson J, Petrauskas L, Atkinson S, Calkins DG (2007) Permanent marking in an endangered species: physiological response to hot branding in Steller sea lions (*Eumetopias jubatus*). Wildl Res 34:43-47
- Merrick RL, Brown R, Calkins DG, Loughlin TR (1995) A comparison of Steller sea lion, *Eumetopias jubatus*, pup masses between rookeries with increasing and decreasing populations. Fish Bull 93:753-758

- Myers MJ, Litz B, Atkinson S (2010) The effects of age, sex, season and geographic region on circulating serum cortisol concentrations in threatened and endangered Steller sea lions (*Eumetopias jubatus*). *Gen Comp Endocrinol* 165:72-77
- Myers MJ, Rea LD, Atkinson S (2006) The effects of age, season and geographic region on thyroid hormones in Steller sea lions (*Eumetopias jubatus*). *Comp Biochem Physiol A* 145:90-98
- Nordoy ES, Aakvaag A, Larsen TS (1993) Metabolic adaptations to fasting in harp seal pups. *Physiol Zool* 66:926-945
- Ortiz RM, Houser DS, Wade CE, Ortiz CL (2003) Hormonal changes associated with the transition between nursing and natural fasting in northern elephant seals (*Mirounga angustirostris*). *Gen Comp Endocrinol* 130:78-83
- Ortiz RM, Noren DP, Litz B, Ortiz CL (2001) A new perspective on adiposity in a naturally obese mammal. *Am J Physiol Endocrinol Metab* 281:E1347-E1351
- Ortiz RM, Wade CE, Ortiz CL (2000) Prolonged fasting increases the response of the Renin-Angiotensin-Aldosterone system, but not vasopressin levels, in postweaned northern elephant seal pups. *Gen Comp Endocrinol* 119:217-223
- Pitcher K, Calkins DG, Pendleton GW (2000) Steller sea lion body condition indices. *Mar Mamm Sci* 16:427-436
- Rea LD (2002) Indices of condition in Steller sea lions (*Eumetopias jubatus*). In: DeMaster D, Atkinson S (eds) *Steller sea lion decline: is it food II*. University of Alaska Sea Grant, Fairbanks, p 29-33
- Romero LM (2002) Seasonal changes in plasma glucocorticoid concentrations in free-living vertebrates. *Gen Comp Endocrinol* 128:1-24
- Rosen DAS, Kumagai S (2008) Hormone changes indicate that winter is a critical period for food shortages in Steller sea lions. *J Comp Physiol B* 178: 573-583
- Sease JL, Taylor WP, Loughlin TR, Pitcher KW (2001) Aerial and land-based surveys of Steller sea lions (*Eumetopias jubatus*) in Alaska, June and July 1999 and 2000. U.S. Dep. Commer., NOAA Tech. Memo. NMFS-AFSC-122, p 52
- Springer AM, Estes JA, van Vliet GB, Williams TM, Doak DF, Danner EM, Forney KA, Pfister B (2003) Sequential megafaunal collapse in the North Pacific Ocean: an ongoing legacy of industrial whaling? *Proc Natl Acad Sci USA* 100:12223-12228

- St. Aubin DJ, Geraci JR (1988) Capture and handling stress suppresses circulating levels of thyroxine (T4) and triiodothyronine (T3) in beluga whales *Delphinapterus leucas*. *Physiol Zool* 61:170-175
- Thomson CA, Geraci JR (1986) Cortisol, aldosterone, and leucocytes in the stress response of bottlenose dolphins, *Tursiops truncatus*. *Can J Fish Aquat Sci* 43:1010-1016
- Trites AW, Jonker RAH (2000) Morphometric measurements and body condition of healthy and starving Steller sea lion pups (*Eumetopias jubatus*). *Aquat Mamm* 26:151-157
- Trites AW, Miller AJ, Alexander MA, Bograd SJ, Calder JA, Capotondi A, Coyle KO, Lorenzo ED, Finney BP, Gregr EJ, Grosch CE, Hare SR, Hunt jr GL, Jahncke J, Kachel NB, Kim H-J, Ladd C, Mantua NJ, Marzban C, Maslowski W, Mendelssohn R, Neilson DJ, Okkonen SR, Overland JE, Reedy-Maschner KL, Royer TC, Schwing FB, Wang JXL, Winship AJ (2007) Bottom-up forcing and the decline of Steller sea lions (*Eumetopias jubatus*) in Alaska: assessing the ocean climate hypothesis. *Fish Oceanogr* 16:46-67
- Trites AW, Porter BT (2002) Attendance patterns of Steller sea lions (*Eumetopias jubatus*) and their young during winter. *J Zool Lond* 256:547-556
- Wilber JF, Utiger RD (1969) The effect of glucocorticoids on thyrotropin secretion. *J Clin Invest* 48:2096-2103

Chapter 4:

Endocrine and immunological responses to adrenocorticotrophic hormone (ACTH) infusion in juvenile harbor seals (*Phoca vitulina*) during winter and spring

4.1 Abstract

Stressors activate the hypothalamic-pituitary-adrenal axis leading to secretion of adrenal hormones including glucocorticoids (GC) and aldosterone. GC have been shown to alter the hypothalamic-pituitary-thyroid axis and the immune system in mammals. Therefore, this study investigated the response of circulating concentrations of several hormones (cortisol, aldosterone, total and free thyroxine and total triiodothyronine) and immunological parameters (total and differential leukocyte counts and peripheral blood mononuclear cells (PBMC) proliferation to an acute stressor, exogenous adrenocorticotrophic hormone (ACTH), during two seasons in juvenile harbor seals. Cortisol concentrations peaked at 165 min in both winter (203.1 ± 84.7 ng/ml) and spring (205.3 ± 65.7 ng/ml) and were still significantly elevated 240 min after ACTH infusion. Aldosterone concentrations peaked at 90 min in both winter (359.3 ± 249.3 pg/ml) and spring (294.1 ± 83.7 pg/ml) and were still elevated 240 min after administration of ACTH in both seasons ($p \leq 0.011$). As in other mammalian species, juvenile harbor seals showed a significant increase in circulating total white blood cells driven primarily by the increase in neutrophil counts, which occurred simultaneously with a decrease in lymphocyte counts leading to an overall decrease in neutrophil to lymphocyte ratio. No clear trend was present in the *in vitro* PBMC proliferation following ACTH administration. There was not a seasonal difference for basal concentration of any hormone quantified or in their response to an acute stressor. Nor was there a seasonal difference in total and differential leukocyte counts or *in vitro* PBMC proliferation. These findings demonstrate that, similar to other mammalian species, an acute stress response in juvenile harbor seals results in an increase in cortisol and aldosterone concentrations, which is associated with altered immunological parameters.

4.2 Introduction

Stress is defined as a state of threatened homeostasis resulting from internal or external environmental change (Stratakis and Chrousos, 1995). The hypothalamic-pituitary-adrenal (HPA) axis is activated in response to a stressor leading to secretion of adrenal hormones including glucocorticoids (GC) and aldosterone (St. Aubin and Geraci, 1986; Thomson and Geraci, 1986; Gulland *et al.*, 1999; Ortiz and Worthy, 2000; Contreras *et al.*, 2004; Romano *et al.*, 2004). In addition to its role in the stress response, cortisol, the prominent GC in pinnipeds (DeRoos and Bern, 1961; Sangalang and Freeman, 1976), regulates gluconeogenesis promoting the mobilization of fatty acids from peripheral adipose tissues. In pinnipeds, cortisol has been suggested to play a role during molt (Riviere *et al.*, 1977; Ashwell-Erickson *et al.*, 1986; Boily, 1996), lactation (Engelhard *et al.*, 2002), fasting (Ortiz *et al.*, 2001a; Ortiz *et al.*, 2001b; Ortiz *et al.*, 2003a; Rosen and Kumagai, 2008) and potentially during diving (Zapol *et al.*, 1979). While aldosterone secretion is influenced by the HPA axis during a stress response, it is predominately under regulation by the renin-angiotensin system in pinnipeds (Malvin *et al.*, 1975; Ortiz *et al.*, 2000; Houser *et al.*, 2001; Ortiz *et al.*, 2003b; Ortiz *et al.*, 2006) with secretion stimulated by angiotensin II in order to maintain water and electrolyte balance in mammals. GC have also been shown to affect the hypothalamic-pituitary-thyroid (HPT) axis (Re *et al.*, 1976; Bianco *et al.*, 1987) and the immune system (Marketon and Glaser, 2008).

Thyroid hormones including total and free thyroxine (T_4) and total triiodothyronine (T_3) are essential for growth and development, metabolism and thermoregulation. Elevated GC concentrations following an acute stressor have been associated with changes in concentrations of T_4 and T_3 . In free ranging beluga whales, T_4 and T_3 concentrations decreased during the 24 hrs following capture and handling (St. Aubin and Geraci, 1988). Similarly, total and free T_4 and total T_3 concentrations in free ranging Steller sea lion pups decreased following the initial rookery disturbance associated with research activities (Chapter 3). In both these studies on free ranging marine mammals basal concentrations of hormones were not known. Therefore, it was

not possible to determine if T_4 and T_3 continually decreased or initially increased followed by a subsequent decrease in response to a stressor. Studies on domesticated animals were able to establish basal concentrations of hormones prior to exposure to an acute stressor. In domestic camels transportation led to an initial increase in cortisol, T_4 and T_3 concentrations followed by a subsequent decrease resulting in these hormones returning to basal concentrations within 24 hr of the acute stressor (Saeb *et al.*, 2010).

Administrations of ACTH or cortisol are associated with characteristic changes in circulating populations of leukocytes and alterations in cell-mediated immune function in cattle and sheep (Roth and Kaeberle, 1981a; Roth and Kaeberle, 1981b). Circulating neutrophil counts increased rapidly with a concurrent decrease in lymphocyte and monocyte counts in association with elevated GC concentrations during restraint in rats (Dhabhar *et al.*, 1995; Dhabhar *et al.*, 1996). Further, suppression of immune function following an acute stressor included reduced *in vitro* proliferation of PBMC in piglets (Westly and Kelley, 1984) but not in lambs (Minton and Blecha, 1990; Coppinger *et al.*, 1991) or calves (Manak, 1986). The differences observed in the effect of elevated GC on immune parameters is dependent on the concentration of GC induced by the acute stressor, the immune parameter and the species being studied (Khansari *et al.*, 1990). Given the physiological influence of GC following exposure to an acute stressor, this study investigated the response of circulating concentrations of several hormones (cortisol, aldosterone, total and free T_4 and T_3) and immunological parameters (total and differential leukocyte count and PBMC proliferation) in harbor seals (*Phoca vitulina*) following administration of exogenous ACTH. I hypothesized that 1) exogenous ACTH would induce an acute stress response as demonstrated by an increase in cortisol concentrations; and 2) the acute stress response would be associated with altered immune parameters including an increase in total WBC and a decrease in *in vitro* PBMC proliferation.

The importance of body mass and composition in harbor seals is evident by the positive correlation of body mass and lipid stores with survival during the first year of independence (Muelbert *et al.*, 2003; Harding *et al.*, 2005). Further, mass of harbor seals

at birth and postnatal growth rates were influenced by the mass of the adult female harbor seal, with heavier females giving birth to heavier pups with a faster growth rate (Ellis *et al.*, 2000; Bowen *et al.*, 2001). Harbor seals also display seasonal variation in body mass, lipid stores, and resting metabolic rate (Rosen and Renouf, 1997; Rosen and Renouf, 1998; Mellish *et al.*, 2007). Given the potential influence of body mass on the health and survival of harbor seals and the seasonality in body mass, we also assessed if body mass was related to an animal's ability to respond to an acute stressor. The objectives of the current study were to determine: (1) the effect of an acute stressor on the HPT axis and immune parameters; (2) if there is a difference in the physiological response to an acute stressor between seasons; and (3) if variation in body mass was associated with variations in the physiological response to an acute stressor as measured by circulating cortisol and aldosterone concentrations.

4.3 Materials and Methods

4.3.1 Animals and Blood Collection

Seven juvenile (1.5 – 3 years old), female harbor seals (*Phoca vitulina*) maintained at the Alaska SeaLife Center in Seward, Alaska (60° N latitude, 149° W longitude) participated in two ACTH administration trials during the winter (November/December 2006) and spring (May 2007). Seals were captured as pups from the declining population in Prince William Sound in 2004 (n=3) and 2005 (n=4). All seals were simultaneously participating in a long-term nutritional study (L. Polasek, personal communication) and were maintained on a mixed diet comprised of pollock (*Theragra chalcogramma*), capelin (*Mallotus villosus*), squid (*Loligo opalescens*) and either Pacific herring (*Clupea pallasii*) or Atlantic herring (*Clupea harengus*). Body mass (BM) was measured to the nearest tenth of a kilogram with a platform scale (Transcell Model TI-500-SL, Accurate Scales, Terre Haute, IN). The seals' mass ranged between 23.5 and 36.4 kg (mean 29.9 ± 1.3 kg) and they were considered healthy based on clinical and behavioral observations by veterinary and animal care staff.

During each trial seals were given a 5 µg/kg of Cortrosyn™ (Amphastar Pharmaceuticals, Inc. Rancho Cucamonga, CA) administered intramuscularly. This dose of Cortrosyn™, a synthetic ACTH, has been previously used in assessment of adrenal function in dogs (Watson *et al.*, 1998; Behrend *et al.*, 2006). Two hours prior to ACTH administration seals were given torbugesic (Fort Dodge; Fort Dodge, Iowa) and diazepam (Hospira Inc.; Lake Forest, Ill) prior to the placement of an indwelling 16 g x 15 cm long-term, guide wire style catheter (Mila International, Inc., Florence, KY, USA) in the extradural vein for serial blood sample collection. Seals were given two hours following catheter placement to recover prior to administration of ACTH. Blood was collected immediately prior to ACTH administration (0 min) and 60, 90, 165, 240 min after ACTH administration. Due to the limited number of seals participating in the study, no saline controls were used and blood collected prior to ACTH administration (time 0) was used for baseline values for all parameters in the present study. Bloods samples (< 17 ml) were drawn at each time point for a total blood volume for each seal ranging between 70 – 85 ml. Cell Preparation Tube (CPT) (CPT™ Vacutainer, Becton Dickinson, Franklin Lakes, NJ) and Vacuette® blood collection tubes were kept upright and chilled until further processing in the lab (< 2 hrs). EDTA-treated blood tubes were collected for total and differential white blood cell (WBC) counts. CPT with sodium citrate was collected for isolation of PBMC. Serum was collected and stored at -80°C until hormone concentrations were quantified.

4.3.2 Hormone Radioimmunoassays

Commercially available radioimmunoassay (RIA) kits were used to quantify hormone concentrations in harbor seal serum. All samples were run in duplicate per manufacturer instructions with the exception that all volumes were halved. Solid-phase RIA kits (Siemens, Los Angeles, CA) were used for quantifying cortisol, aldosterone, total and free T₄ and total T₃ concentrations. Cortisol, total and free T₄ and total T₃ were previously validated for use with harbor seal serum in our laboratory (Oki and Atkinson, 2004) and aldosterone was validated in the present study. Pools of male and female

harbor seal serum were serially diluted to determine degree of parallelism to the standard curve. Recovery of added aldosterone (25–1200 pg/ml) for males ($y = -21.591x + 173.63$, $r^2 = 0.996$) and females ($y = -23.63x + 282.29$; $r^2 = 0.998$) was determined. Inter-assay and intra-assay coefficients of variation were $< 10.0\%$ for each hormone. Assay sensitivities were as follows: cortisol 2.0 ng/ml, aldosterone 11.0 pg/ml, total T_4 2.5 ng/ml, free T_4 0.1 pg/ml, and total T_3 0.07 ng/ml.

4.3.3. Total and Differential Leukocyte Counts

Total WBC counts were determined using the CBC-Diff Veterinary Hematological System (Heska ® Corporation, Loveland, CO). Blood smears were made from whole EDTA blood and stained with Wright-Giemsa (Dip Quick Stain, Jorgensen Laboratories, Loveland CO). WBC differentials, including neutrophils, lymphocytes, monocytes and eosinophils were counted manually and are reported as both percentages and absolute values of differentials.

4.3.4 PBMC Isolation and Proliferation

Blood collected into CPT tubes was processed per manufacturer instructions. Briefly, CPT tubes were centrifuged at $1500 \times g$ for 20 min and the PBMC were re-suspended in freezing media comprised of RPMI 1640, 20% FBS (Hyclone, Thermo Fisher Scientific Inc., Logan, UT), 10% DMSO (Sigma Chemical Co., St. Louis, MO) and stored in liquid nitrogen until analysis. Cryopreserved PBMC were thawed and washed twice with PBS (without Ca^{2+}/Mg^{2+}) and re-suspended in RPMI 1640 supplemented with 10% FBS, 2 mM L-glutamine, penicillin (100 U/ml), and streptomycin (100 μ g/ml), 1 mM sodium pyruvate, 100 μ M nonessential amino acids, 25 mM hepes, and 0.05 mM 2-mercaptoethanol (Sigma Chemical Co., St. Louis, MO). All cell culture media and supplements, except for FBS, DMSO and 2-mercaptoethanol, were purchased from Gibco, Invitrogen Corp., Carlsbad, CA. Viability was $>90\%$ and assessed using the exclusion dye trypan blue (Sigma Chemical Co., St. Louis, MO).

PBMC were plated (2.0×10^5 cells/well) in 96-well flat-bottom tissue-culture plates (Falcon, Becton Dickinson, Franklin Lakes, NJ) and their proliferation assessed following exposure to either concanavalin A (ConA, Sigma Aldrich, St. Louis, MO) or lipopolysaccharide (LPS055:B5, Sigma Aldrich, St. Louis, MO). PBMC were cultured at 37°C with 5% CO₂ in triplicate for each time point for all seven harbor seals in each of the following treatment groups; 1) unexposed control, 2) ConA suboptimal (0.1 µg/ml), 3) ConA optimal (1.0 µg/ml), 4) LPS suboptimal (1.0 µg/ml) and 5) LPS optimal (100 µg/ml). ConA is a plant-derived mitogen and preferentially stimulates T cells (Barta and Barta, 1993) while LPS from *Escherichia coli* is a B-cell activator (Wechsler-Reya and Monroe, 1996). The use of suboptimal and optimal mitogen concentrations have been used in previous pinniped studies (Levin *et al.*, 2005; Mori *et al.*, 2006; Keogh *et al.*, 2010). Mitogens were reconstituted in RPMI 1640 supplemented with 10% FBS, 2 mM L-glutamine, penicillin (100 U/ml), and streptomycin (100 µg/ml). These conditions are similar to those previously used for pinnipeds (Levin *et al.*, 2005; Mori *et al.*, 2006; Keogh *et al.*, 2010).

All PBMC were assessed in two assays, one for each season, with PBMC collected from a captive adult Steller sea lion (*Eumetopias jubatus*) as an inter-assay control. PBMC were incubated for 48 hrs after which 20 µl of bromodioxymuridine (BrdU) was added for a final concentration of 10 µM BrdU and incubated for an additional 18 hrs for a total incubation period of 66 hrs. Proliferation was assessed with the Cell Proliferation Biotrack ELISA System version 2 (GE Healthcare, Piscataway, NJ) per manufacturer's instructions using Dynatech MRX Revelation microplate reader (Dynatech Laboratories, Inc, Chantilly, VA) at 450 nm with a reference wavelength of 650 nm. Data are presented as mean of the triplicates for optical density (OD) with standard error and as a stimulation index (SI; mean OD of cells exposed to mitogen/mean OD of cells in media only).

4.3.5 Statistical Analysis

Data were analyzed with Systat 10 (Systat Software, Inc, Point Richmond, CA). Data were analyzed with Systat 10 (Systat Software, Inc., Point Richmond, CA, USA). For the purposes of analyses ACTH trials conducted in November or December of 2006 were defined as winter while spring trials took place during May 2007. Repeated measures analysis of variance (ANOVA) was used to evaluate the influence of season for each parameter. Additionally, the area under the curve was calculated based on the trapezoid formula with reference to the baseline concentrations (AUC_I) for cortisol and aldosterone concentrations following administration of ACTH (Pruessner *et al.*, 2003; Fekedulegn *et al.*, 2007). Paired t-tests were used to determine differences between baseline values at 0 min and the AUC_I between seasons. Pearson correlations were used to assess the association of body mass with basal hormone concentrations, total and differential WBC counts and *in vitro* proliferation of PBMC within each season. Results were considered statistically significant at $p \leq 0.050$.

4.4 Results

4.4.1 Changes in Hormone Concentrations Following ACTH Administration

Basal (0 min) serum cortisol concentrations did not differ (Table 4.1, $p=0.461$) between winter (28.5 ± 10.3 ng/ml) and spring (35.8 ± 21.6 ng/ml). Nor did basal aldosterone concentrations differ ($p=0.472$) between winter (18.2 ± 8.4 pg/ml) and spring (38.1 ± 29.6 pg/ml). In both seasons cortisol ($p \leq 0.001$) and aldosterone ($p \leq 0.001$) significantly increased in response to ACTH administration (Figure 4.1). Circulating concentrations of cortisol (winter: $p < 0.001$; spring: $p = 0.001$) and aldosterone (winter: $p = 0.003$; spring: $p < 0.001$) were significantly increased 60 min after administration of ACTH compared to basal concentrations in all seals and in both seasons (Figure 4.1). Cortisol concentrations peaked at 165 min in both winter (203.1 ± 84.7 ng/ml) and spring (205.3 ± 65.7 ng/ml) and were still significantly elevated 240 min after ACTH administration in both seasons ($p \leq 0.001$). Aldosterone concentrations peaked at 90 min in both winter (359.3 ± 249.3 pg/ml) and spring (294.1 ± 83.7 pg/ml) and were still elevated

240 min after administration of ACTH in both seasons ($p \leq 0.011$). There was no significant difference between seasons in cortisol ($p=0.872$) or aldosterone ($p=0.732$) concentrations in response to ACTH administration (Figure 4.1). Similarly, there was no difference between AUC₁ between seasons for cortisol ($p=0.287$) or aldosterone ($p=0.326$) (Figure 4.1). In both seasons aldosterone and cortisol concentrations were positively correlated ($p \leq 0.003$).

There was a large amount of individual variation observed in the response of thyroid hormones to ACTH administration (Figure 4.2). Circulating concentrations of total T₄ ($p=0.032$) and total T₃ ($p=0.011$) but not free T₄ ($p=0.522$) significantly changed in response to ACTH stimulation (Figure 4.2). There was not a seasonal difference in the response to ACTH stimulation for total T₄ ($p=0.116$), free T₄ ($p=0.366$), nor total T₃ ($p=0.861$).

There was no difference between winter and spring for body mass (winter 29.4 ± 5.3 kg; spring 30.5 ± 4.6 kg; $p=0.184$). Basal cortisol concentrations were correlated to body mass in winter ($p=0.039$) but not in the spring ($p=0.311$). Further, there was no relationship between mass and cortisol concentrations at any time following ACTH administration in winter ($p<0.087$). In spring, response in cortisol to ACTH administration was related to body mass ($p=0.046$). Basal aldosterone concentrations were not correlated to body mass in either season ($p>0.154$). Further, there was no relationship between body mass and aldosterone concentrations at any time following ACTH administration in either season ($p<0.112$).

Table 4.1. Basal (time 0) mean (\pm SD) and range of cortisol, aldosterone, total and differential blood cell counts and PBMC proliferation by season from captive juvenile harbor seals (n=7).

Parameter	Winter				Spring			
	Mean \pm SD		Range		Mean \pm SD		Range	
Cortisol (ng/ml)	28.5 \pm	10.3	13.2	– 46.7	35.8 \pm	21.6	18.0	– 75.4
Aldosterone (pg/ml)	18.2 \pm	8.4	11.1	– 35.1	30.8 \pm	29.6	4.9	– 72.0
Total thyroxine (ng/ml)	26.8 \pm	5.2	18.6	– 33.5	21.9 \pm	5.0	13.9	– 28.3
Free thyroxine (pg/ml)	6.4 \pm	2.3	3.6	– 9.9	4.9 \pm	1.4	3.2	– 7.3
Total triiodothyronine (ng/ml)	0.7 \pm	0.2	0.5	– 0.1	0.74 \pm	0.1	0.6	– 0.9
WBC ($10^3/\mu$ l)	8.6 \pm	1.5	6.7	– 11.0	7.9 \pm	1.5	6.3	– 10.0
Neutrophil ($10^3/\mu$ l)	5.4 \pm	0.7	4.7	– 6.5	5.0 \pm	1.3	2.8	– 7.1
(% of WBC)	66.0 \pm	8.3	56	– 72	63.1 \pm	7.5	54	– 72
Lymphocyte ($10^3/\mu$ l)	1.9 \pm	0.7	1.1	– 2.9	2.1 \pm	0.7	1.6	– 3.4
(% of WBC)	22.4 \pm	4.7	17	– 29	26.3 \pm	5.2	7	– 20
Monocytes ($10^3/\mu$ l)	0.7 \pm	0.3	0.3	– 1.1	0.5 \pm	0.3	0.1	– 0.9
(% of WBC)	8.9 \pm	3.1	4	– 14	5.7 \pm	2.6	1	– 9
Eosinophil ($10^3/\mu$ l)	0.2 \pm	0.2	0.0	– 0.5	0.1 \pm	0.1	0.1	– 0.3
(% of WBC)	2.0 \pm	2.3	0	– 5	1.7 \pm	0.7	1	– 3
N:L ratio	3.1 \pm	1.0	1.9	– 4.3	2.6 \pm	0.9	1.6	– 4.0
Control (OD)	0.29 \pm	0.02	0.25	– 0.33	0.30 \pm	0.05	0.23	– 0.36
ConA 0.1 μ g/ml (OD)	0.29 \pm	0.01	0.26	– 0.30	0.31 \pm	0.05	0.29	– 0.39
ConA 0.1 μ g/ml (SI)	1.01 \pm	0.07	0.88	– 1.10	1.06 \pm	0.08	0.91	– 1.19
ConA 1.0 μ g/ml (OD)	1.57 \pm	0.58	0.29	– 1.95	2.00 \pm	0.30	1.69	– 2.63
ConA 1.0 μ g/ml (SI)	5.55 \pm	2.18	0.95	– 7.50	6.87 \pm	1.12	5.22	– 8.81
LPS 055:B5 1.0 μ g/ml (OD)	0.37 \pm	0.09	0.29	– 0.53	0.58 \pm	0.19	0.41	– 0.83
LPS 055:B5 1.0 μ g/ml (SI)	1.27 \pm	0.23	0.94	– 0.63	1.51 \pm	0.39	0.94	– 1.63
LPS 055:B5 100 μ g/ml (OD)	0.68 \pm	0.28	0.28	– 1.01	0.88 \pm	0.35	0.53	– 1.60
LPS 055:B5 100 μ g/ml (SI)	2.38 \pm	1.04	0.91	– 3.97	2.91 \pm	0.77	2.10	– 4.50




Figure 4.1. Mean (\pm SD) serum cortisol (A) and aldosterone (C) concentrations over time following ACTH injection during the winter (●) and spring (○) for seven juvenile harbor seals ($n=7$). Area under the curve (AUC_I) following ACTH injection for cortisol (B) and aldosterone (D) during the winter and spring experiments. * denotes significant ($p < 0.05$) difference from time 0.

4.4.2 Alterations in Total and Differential Leukocyte Counts

There was no difference in response to ACTH administration between seasons in circulating total WBC count (Figure 4.3, $p=0.241$). Nor was there a difference in neutrophils, monocytes or eosinophils whether expressed as percentage or absolute number ($p>0.062$), or neutrophil: lymphocyte (N: L) ratio ($p<0.142$). There was a significant effect of season on circulating lymphocytes in response to ACTH administration as a percentage ($p=0.020$) with winter having lower percent of lymphocytes compared to spring. This relationship was not found when lymphocytes were reported as an absolute count ($p=0.537$). Total WBC count ($p<0.001$), neutrophils (percentage, $p<0.001$; absolute count, $p<0.001$), and lymphocytes (percentage, $p\leq 0.001$; absolute count, $p<0.001$) changed following ACTH administration (Figure 4.3). However, no change was observed in circulating monocytes (percentage, $p=0.229$; absolute count, $p=0.386$). Specifically, total WBC, neutrophil counts and N: L ratios were significantly elevated 240 min after ACTH administration ($p<0.021$) while lymphocyte count decreased ($p<0.007$). Further, there was no relationship between total or differential WBC count and between body mass for either season ($p>0.141$).

4.4.3 PBMC *in vitro* Proliferation

There was no difference in response to ACTH administration between seasons in the spontaneous (unexposed) or stimulated (ConA or LPS) PBMC proliferation ($p>0.089$). Nor did spontaneous ($p=0.830$) or suboptimal ConA (OD, $p=0.243$; SI, $p=0.752$) PBMC proliferation change following ACTH administration. There was no difference in PBMC proliferation between seasons when stimulated by optimal ConA but proliferation did significantly change following ACTH administration (OD, $p<0.001$; SI, $p<0.001$); however, while there was a clear pattern, there was not an overall trend present (Figure 4.4). PBMC-proliferation did significantly change following ACTH administration when stimulated by suboptimal (OD, $p=0.006$; SI, $p=0.036$) and optimal LPS (OD $p=0.001$; SI $p=0.005$). Similar to the response in stimulation by optimal ConA, proliferation following *in vitro* exposure to LPS (optimal and suboptimal) showed a clear pattern but with no obvious trend. *In vitro* PBMC proliferation was only positively correlated with body

mass following stimulation with optimal LPS at time 0 in the winter when reported as an OD ($p=0.046$) but not SI ($p=0.120$).




Figure 4.2. Mean (\pm SD) serum concentration for total (A) and free (B) thyroxine and total triiodothyronine (C) following ACTH injection during the winter (●) and spring (○) for seven juvenile harbor seals ($n=7$). Dashed lines denote baseline concentrations.




Figure 4.3 Mean differential leukocyte counts for all seven juvenile harbor seals following an acute stressor (exogenous ACTH) during the winter with results reported as percentage (A) and absolute count (B) and during spring with results reported as percentage (C) and absolute count (D).




Figure 4.4 PBMC proliferation responses from seven captive juvenile harbor seals exposed to (A) 1.0 µg/ml ConA and (B) 100 µg/ml LPS following an acute stressor during winter (●) and spring (○). Results presented as mean stimulation index (SI) ± SD.

4.5 Discussion

4.5.1 Hormone Concentrations Induced by ACTH

An acute stress response was induced in all seals as evident by the changes in cortisol concentrations following administration of 5 µg/kg of Cortrosyn™. Gulland *et al.*, (1999) reported cortisol concentrations prior to ACTH administration higher in harbor seal pups (2-4 months old) during rehabilitation than basal cortisol concentrations in the present study. Similarly, basal cortisol concentrations previously reported for wild and captive harbor seals (Gardiner and Hall, 1997) were higher than concentrations in present study. Peak cortisol concentrations following exogenous ACTH gel (Gulland *et al.*, 1999) resulted in similar peak concentrations as in the present study; however, peak cortisol were observed later (165 min) following ACTH administration in the present study than in the previous study (90 min) (Gulland *et al.*, 1999). These results may be explained by differences in the methodologies employed in the studies. In the present study, an indwelling catheter was used for serial blood collection, following the

placement of which seals were allowed to rest for 2 hrs prior to administration of ACTH. This experimental set up eliminated the need to physically restrain the seals during blood collection and allowed seals to recover from the increase in cortisol concentrations measured at the time of the catheter placement (data not shown). During the 240 min following ACTH administration in the present study, seals remained calm and most slept for large periods of time. It is important to note that the cortisol concentrations observed in the present study including the peak observed at 165 min following ACTH administration were lower than reported concentrations of cortisol in wild harbor seals (Gardiner and Hall, 1997; Gulland *et al.*, 1999). This finding was not unexpected given the differences between harbor seal populations and since different stressors result in different behavioral and physiological responses including cortisol concentrations. The increase in cortisol concentrations differed based on capture methods used in bears (Cattet *et al.*, 2003), white-tailed deer (Delgiudice *et al.*, 1990; Denicola and Swihart, 1997), bottlenose dolphins (Thomson and Geraci, 1986) and elephant seals (Engelhard *et al.*, 2002).

While cortisol concentrations have been frequently reported for harbor seals in relation to molt (Riviere *et al.*, 1977; Ashwell-Erickson *et al.*, 1986), breeding (Gardiner and Hall, 1997), seasonal and diurnal patterns (Gardiner and Hall, 1997; Oki and Atkinson, 2004) there is comparatively limited information available for aldosterone concentrations in harbor seals (St. Aubin and Geraci, 1986; Gulland *et al.*, 1999). The present study showed a significant increase in aldosterone concentration, which was correlated with cortisol concentrations following ACTH administration, demonstrating that secretion of aldosterone is also under the HPA axis control as has been previously demonstrated in phocids (St. Aubin and Geraci, 1986; Gulland *et al.*, 1999). Basal aldosterone concentrations or concentrations in response to ACTH administration were not related to season. Similar to cortisol, basal aldosterone concentrations in the present study were lower during both seasons than previous reported initial aldosterone concentrations in harbor seals pups during rehabilitation or wild harbor seals (Gulland *et al.*, 1999). Gulland *et al.*, (1999) reported a peak in aldosterone concentrations 90 min

following ACTH administration, similar to the present study; however, harbor seal pups given an ACTH administration during rehabilitation had more than a 4 times greater aldosterone concentrations compared to the 90 min peak in the present study. Given that aldosterone secretion is under the control of HPA as well as renin-angiotensin system, the higher aldosterone concentrations previously reported for harbor seals (Gulland *et al.*, 1999) may be due to disease and physiological differences between study populations and methodologies such as different ACTH and restraint during blood collections.

Basal cortisol concentrations were correlated to body mass in winter but not spring, which may indicate that body mass is more critical during the winter. Similar to the lack of seasonal differences in body mass in the present study we found no seasonal difference in cortisol or aldosterone concentrations in response to exogenous ACTH. While these findings may be, at least in part, due to the seals being immature, captive seals have been reported to lack seasonal variation in cortisol concentrations observed in wild seals (Gardiner and Hall, 1997). Gardiner and Hall (1997) found circulating cortisol concentrations in mature captive females did not show the seasonal variation observed in mature wild female and male harbor seals, which was attributed to the consistent environment and availability of food in the captive environment.

4.5.2 Alterations in Total and Differential Leukocyte Counts

The basal WBC count in the present study was similar to previously published results for captive harbor seals (McConnell and Vaughan, 1983). Compared to wild adult harbor seals (Ross *et al.*, 1993; Morgan *et al.*, 1998) baseline WBC, neutrophil, and eosinophil counts were lower while lymphocyte and monocytes counts were higher in the present study. Differences in total and differential WBC counts may be indicative of population difference. However, differences in total and differential white blood cell counts between immature and mature mammals have been reported across taxa including carnivore (Nunn *et al.*, 2009). Therefore, the differences in total and differential leukocyte counts observed in the present study may be attributed to age, since Nielsen (1995) found neutrophils (%) increased while lymphocytes (%) decreased with age. The

lower eosinophil counts observed in the present study as in other captive seal populations are likely due to the expected lower occurrence of parasites in captive compared to wild animals. In the present study total and differential WBC counts were not related to body mass. Across mammalian species the number of circulating total white blood cell counts increased with body mass in adults (Nunn *et al.*, 2003; Nunn *et al.*, 2009). However, the present study did not find body mass to be related to total WBC count.

The immune system is not independent but rather there is a well-established link between the immune and endocrine systems (Haddad *et al.*, 2002; Weber, 2003). Capture and handling methods routinely used during blood collection in wild mammals can result in elevated cortisol concentrations (Engelhard *et al.*, 2002; Romero *et al.*, 2008; Keogh *et al.*, 2010), which may alter hematological parameters (Castellini *et al.*, 1996; Cattet *et al.*, 2003). Circulating neutrophil counts increase rapidly with a concurrent decrease in lymphocyte and monocyte counts during an acute stress response in association with elevated GC concentrations in terrestrial mammals (Dhabhar *et al.*, 1995; Dhabhar *et al.*, 1996; Cattet *et al.*, 2003). Similarly, an increase in total WBC, neutrophil and neutrophil to lymphocyte ratio was associated with increases in cortisol concentration following ACTH administration in the present study. The observed increase in total WBC and neutrophil to lymphocyte ratio was driven predominately by the increase in neutrophils. Juvenile harbor seals response to an acute stressor in the present study was typically mammalian exhibiting a stress leukogram.

4.5.3 PBMC *in vitro* Proliferation

PBMC including lymphocytes play an important role in cell-mediated immunity. The immunosuppressive effect of elevated GC concentration associated with stressors has been well established in mammals (Keller *et al.*, 1981; Deguchi and Akuzawa, 1998; Bilandzic *et al.*, 2006). Elevated cortisol concentrations have been shown to suppress PBMC proliferation in piglets (Westly and Kelley, 1984) while having no influence on lambs (Minton and Blecha, 1990) or calves (Manak, 1986). The effect of elevated GC on immune parameters is dependent on several variables including the concentration of GC

induced by an acute stressor, the immune parameter or function, and the species being studied (Bilandzic *et al.*, 2006). Previous studies on the acute stress response in seals (St. Aubin and Geraci, 1986; Gulland *et al.*, 1999) have not assessed the impact of an acute stressor on total and differential leukocyte count or *in vitro* PBMC proliferation. Therefore, we assessed the effect of an acute stressor (exogenous ACTH) on *in vitro* PBMC proliferation in addition to total and differential leukocyte counts. While we did not quantify the number of circulating T and B cells, the use of two mitogens that preferentially stimulate either T cells (Barta and Barta, 1993) or B cells (Wechsler-Reya and Monroe, 1996) allowed the assessment of both cell populations. All harbor seals in the present study responded to *in vitro* stimulation with ConA and LPS 055:B5 indicating peripheral T and B cells are capable of responding to an antigenic challenge as previously reported for harbor seals (Ross *et al.*, 1993; Swart *et al.*, 1993; Levin *et al.*, 2005). As with previous studies which assessed PBMC proliferation in harbor seals, T cells proliferated more so than B cells when exposed to *in vitro* mitogens at optimal and suboptimal concentrations (Swart *et al.*, 1993; Levin *et al.*, 2005). The lower B cell proliferation in these studies may be the result of the low number of B cells in the peripheral blood as observed in other mammals (Byrne *et al.*, 2000; Faldyna *et al.*, 2001). However, the ratio of T and B cells in harbor seals will need to be confirmed in future studies. T cell proliferation in the present study, regardless of time after ACTH administration was higher than reported for wild lactating female and harbor seal pups (Ross *et al.*, 1993). The SI for B cell proliferation were within the range previously reported for harbor seals (Levin *et al.*, 2005). Levin *et al.*, (2005) reported a negative relationship between B cell proliferation (SI) and body mass in harbor seal pups which was not observed in the present study. However, B cells proliferation in the present study was related to body mass in the winter for optimal LPS when reported as OD but this relationship was not maintained when proliferation was reported as SI.

4.5.4 Body Mass

Body mass has been shown to be important in the health and survival of harbor seals (Bowen *et al.*, 2001; Muelbert *et al.*, 2003; Harding *et al.*, 2005). Body mass is influenced by an individual's disease and nutritional state; therefore, we assessed the influence of body mass on the seal's response to an acute stressor. Harbor seals have been observed to first give birth between 4 and 6 years of age (Markussen *et al.*, 1989), therefore the female harbor seals (1.5 – 3 years old) in the present study were likely immature. As expected, seals in the present study had lower body mass compared to mature seals (Ellis *et al.*, 2000; Bowen *et al.*, 2001; Mellish *et al.*, 2007). The juvenile harbor seals in the present study showed no seasonal variation in total body mass and had a lower body mass in both seasons than total body mass reported for wild yearling (body mass, 33.3 ± 1.3 kg) and juvenile (44-61 kg) harbor seals (Swart *et al.*, 1994; Muelbert *et al.*, 2003). Mature harbor seals have been reported to display seasonal variation in body mass, lipid stores (Rosen and Renouf, 1997; Mellish *et al.*, 2007), energy intake (Rosen and Renouf, 1998), and resting metabolic rate (Rosen and Renouf, 1998). However, seasonal variation was not found in body mass, energy intake or blubber stores in previous studies on juvenile harbor seals (Pitcher, 1986; Markussen and Ryg, 1990). Therefore, it may be that the lack of seasonal variation in body mass in the present study is due, at least in part, to the seals being immature.

4.6 Conclusions

Juvenile harbor seals did not show seasonal variation in any parameter measured, nor did body mass affect the response to an acute stressor in the present study. However, basal cortisol concentrations were related to body mass in the winter, which is suggestive that body mass in the winter may be important for pinnipeds as suggested by Rosen and Kumagai (2008). Circulating concentrations of cortisol and aldosterone significantly increased in response to an acute stressor (exogenous ACTH), which resulted in a stress leukogram at 240 min after application of the stressor. As in other mammalian species, juvenile harbor seals showed a significant increase in total WBC count driven primarily

by the increase in neutrophil counts, which occurred simultaneously with a decrease in lymphocyte counts leading to an overall decrease in N: L ratio. No clear trend was apparent in the *in vitro* proliferation of T or B cells following ACTH administration.

4.7 Acknowledgements

This research was conducted in partial fulfillment of the requirements for the degree of Doctorate of Philosophy at the University of Alaska, Fairbanks. Funding for this project was provided by the Alaska SeaLife Center with congressionally appropriated funds provided by National Marine Fisheries Service and a grant awarded to S. Atkinson by the Pollock Conservation Cooperative Research Center. However, the findings and conclusions presented by the authors are their own and do not necessarily reflect the views or positions of the Center or the University of Alaska. Additional funding was provided through a fellowship from the National Science Foundation GK-12 award to the University of Alaska Fairbanks “New GK-12 Program: The CASE (Changing Alaska Science Education) for Enhancing Understanding of Climate Change” to M. Keogh. This research was authorized under US Marine Mammal Protection Act Permit 881-1710 with all protocols reviewed and approved by the Institutional Animal Care and Use Committee at ASLC (#03-009). We would like to thank the staff at the ASLC and the University of Alaska Fairbanks School of Fisheries and Ocean Sciences Endocrine Lab. Drs. M. Castellini, T. Hollmen, R.M. Ortiz, J. Runstadler graciously provided comments on this manuscript.

4.8 Literature Cited

- Ashwell-Erickson, S., Fay, F. H., Elsner, R., 1986. Metabolic and hormonal correlates of molting and regeneration of pelage in Alaskan harbor and spotted seals (*Phoca vitulina* and *Phoca largha*). *Can. J. Zool.* 64, 1086-1094.
- Barta, O., Barta, V., 1993. Lymphocyte transformation test. In: Barta, O. (Ed.), *Veterinary Clinical Immunology Laboratory. Monographs in Animal Immunology*, 2. Bar-Lab, Inc Blacksburg, pp. B4-1-B4-17.

- Behrend, E. N., Kemppainen, R. J., Bruyette, D. S., Busch, K. A., 2006. Intramuscular administration of a low dose of ACTH for ACTH stimulation testing in dogs. *J. Am. Vet. Med. Assoc.* 229, 528-530.
- Bianco, A. C., Nunes, M. T., Hell, N. S., Maciel, R. M. B., 1987. The role of glucocorticoids in the stress-induced reduction of extrathyroidal 3,5,3'-triiodothyronine generation in rats. *Endocrinol.* 120, 1033-1038.
- Bilandzic, N., Zuric, M., Lojkic, M., Simic, B., Milic, D., Bara, I., 2006. Cortisol and immune measures in boars exposed to three-day administration of exogenous adrenocorticotrophic hormone. *Vet. Res. Commun.* 30, 433-444.
- Boily, P., 1996. Metabolic and hormonal changes during the molt of captive gray seals (*Halichoerus grypus*). *Am. J. Physiol. Regul. Integr. Comp. Physiol.* 270, 1051-1058.
- Bowen, W. D., Ellis, S. L., Iverson, S. J., Boness, D. J., 2001. Maternal effects on offspring growth rate and weaning mass in harbour seals. *Can. J. Zool.* 79, 1088-1101.
- Byrne, K. M., Kim, H. W., Chew, B. P., Reinhart, G. A., Hayek, M. G., 2000. A standardized gating technique for the generation of flow cytometry data for normal canine and normal feline blood lymphocytes. *Vet. Immunol. Immunopathol.* 73, 167-182.
- Castellini, J. M., Melselman, H. J., Castellini, M. A., 1996. Understanding and interpreting hematocrit measurements in pinnipeds. *Mar. Mamm. Sci.* 12, 251-264.
- Cattet, M. R. L., Christison, K., Caulkett, N. A., Stenhouse, G. B., 2003. Physiologic responses of grizzly bears to different methods of capture. *J. Wildl. Dis.* 39, 649-654.
- Contreras, L. N., Arregger, A. L., Persi, G. G., Gonzalez, N. S., Cardoso, E. M., 2004. A new less-invasive and more informative low-dose ACTH test: salivary steroids in response to intramuscular corticotrophin. *Clin. Endocrinol.* 61, 675-682.
- Coppinger, T. R., Minton, J. E., Reddy, P. G., Blecha, F., 1991. Repeated restraint and isolation stress in lambs increases pituitary-adrenal secretions and reduces cell-mediated immunity. *J. Anim. Sci.* 69, 2808-2814.
- Deguchi, E., Akuzawa, M., 1998. Effects of fighting after grouping on plasma concentration and lymphocyte blastogenesis of peripheral blood mononuclear cells induced by mitogens in piglets. *J. Vet. Med. Sci.* 60, 149-153.

- Delgiudice, G. D., Kunkel, K. E., Mech, L. D., Seal, U. S., 1990. Minimizing capture-related stress on white-tailed deer with a capture collar. *J. Wildl. Manag.* 54, 299-303.
- Denicola, A. J., Swihart, R. K., 1997. Capture-induced stress in white-tailed deer. *Wildl. Soc. Bull.* 25, 500-503.
- DeRoos, C. C., Bern, H. A., 1961. The corticoids of the adrenal of the California sea lion (*Zalophus californianus*). *Gen. Comp. Endocrinol.* 1, 275-285.
- Dhabhar, F. S., Miller, A. H., McEwen, B. S., Spencer, R. L., 1995. Effects of stress on immune cell distribution. *J. Immunol.* 154, 5511-5527.
- Dhabhar, F. S., Miller, A. H., McEwen, B. S., Spencer, R. L., 1996. Stress-induced changes in blood leukocytes distribution: Role of adrenal steroid hormones. *J. Immunol.* 157, 1638-1644.
- Ellis, S. L., Bowen, W. D., Boness, D. J., Iverson, S. J., 2000. Maternal effects on offspring mass and stage of development at birth in the harbor seal, *Phoca vitulina*. *J. Mammal.* 81, 1143-1156.
- Engelhard, G. H., Brasseur, S. M. J. M., Hall, A. J., Burton, H. R., Reijnders, P. J. H., 2002. Adrenocortical responsiveness in southern elephant seal mothers and pups during lactation and the effect of scientific handling. *J. Comp. Physiol. B.* 172, 315-328.
- Faldyna, M., Leva, L., Knotigova, P., Toman, M., 2001. Lymphocyte subsets in peripheral blood of dogs - a flow cytometric study. *Vet. Immunol. Immunopathol.* 82, 23-37.
- Fekedulegn, D. B., Andrew, M. E., Burchfiel, C. M., Hartley, T. A., Charles, L. E., Miller, D. B., 2007. Area under the curve and other summary indicators of repeated waking cortisol measurements. *Psychosom. Med.* 69, 651-659.
- Gardiner, K. J., Hall, A. J., 1997. Diel and annual variation in plasma cortisol concentrations among wild and captive harbor seals (*Phoca vitulina*). *Can. J. Zool.* 75, 1773-1780.
- Gulland, F. M. D., Haulena, M., Lowenstine, L. J., Munro, C., Graham, P. A., Bauman, J., Harvey, J., 1999. Adrenal function in wild and rehabilitated pacific harbor seals (*Phoca vitulina richardii*) and in seals with phocine herpesvirus-associated adrenal necrosis. *Mar. Mamm. Sci.* 15, 810-827.

- Haddad, J. J., Saade, N. E., Safieh-Garabedian, B., 2002. Cytokines and neuro-immune-endocrine interactions: a role for the hypothalamic-pituitary-adrenal revolving axis. *J. Neuroimmunol.* 133, 1-19.
- Harding, K. C., Fugiwara, M., Axberg, Y., Harkonen, T., 2005. Mass-dependent energetics and survival in harbour seal pups. *Func. Ecol.* 19, 129-135.
- Houser, D. S., Crocker, D. E., Webb, P. M., Costa, D. P., 2001. Renal function in suckling and fasting pups of the northern elephant seal. *Comp. Biochem. Physiol. A: Mol. Integr. Physiol.* 129, 405-415.
- Keller, S. E., Weiss, J. M., Schleifer, S. J., Miller, N. E., Stein, M., 1981. Suppression of immunity by stress: effect of a graded series of stressors on lymphocyte stimulation in the rat. *Science.* 213, 1397-1400.
- Keogh, M. J., Maniscalco, J. M., Atkinson, S., 2010. Steller sea lion (*Eumetopias jubatus*) pups undergo a decrease in circulating white blood cells and the ability of T cells to proliferate during early postnatal development. *Vet. Immunol. Immunopathol.* 137, 298-304.
- Khansari, D. N., Murgo, A. J., Faith, R. E., 1990. Effects of stress on the immune system. *Immunol. Today* 11, 170-175.
- Levin, M., De Guise, S. D., Ross, P. S., 2005. Association between lymphocyte proliferation and polychlorinated biphenyls in free-ranging harbor seal (*Phoca vitulina*) pups from British Columbia, Canada. *Environ. Toxicol. Chem.* 24, 1247-1252.
- Malvin, R. L., Ridgway, S. H., Cornell, L., 1975. Renin and aldosterone levels in dolphins and sea lions (40117). *Proc. Soc. Exp. Biol. Med.* 157, 665-668.
- Manak, R. C., 1986. Neonatal calf serum immunoregulation of lymphocyte blastogenic responses. *J. Anim. Sci.* 62, 759-771.
- Marketon, J. I. W., Glaser, R., 2008. Stress hormones and immune function. *Cell. Immunol.* 252, 16-26.
- Markussen, N. H., Bjorge, A., Oritsland, N. A., 1989. Growth in harbour seals (*Phoca vitulina*) on the Norwegian coast. *J. Zool. Lond.* 219, 443-440.
- Markussen, N. H., Ryg, M., 1990. Energy requirements for maintenance and growth of captive harbour seals, *Phoca vitulina*. *Can. J. Zool.* 68, 423-426.

- McConnell, L. C., Vaughan, R. W., 1983. Some blood values in captive and free-living common seals (*Phoca vitulina*). *Aquat. Mamm.* 10, 9-13.
- Mellish, J.-A. E., Horning, M., York, A. E., 2007. Seasonal and spatial blubber depth changes in captive harbor seals (*Phoca vitulina*) and Steller's sea lions (*Eumetopias jubatus*). *J. Mammol.* 88, 408-414.
- Minton, J. E., Blecha, F., 1990. Effect of acute stressors on endocrinological and immunological functions in lambs. *J. Anim. Sci.* 68, 3145-3151.
- Morgan, L., Kumaresan, S., Thomas, C., MacWilliams, P., 1998. Hematology and chemistry reference values for free-ranging harbor seals (*Phoca vitulina*) and the effects of hemolysis on chemistry values of captive harbor seals. *J. Zoo Wildl. Med.* 29, 394-400.
- Mori, C., Morsey, B., Levin, M., Nambiar, P. R., De Guise, S. D., 2006. Immunomodulatory effects of *in vitro* exposure to organochlorines on T-cell proliferation in marine mammals and mice. *J. Toxicol. Environ. Health A.* 69, 283-302.
- Muelbert, M. M., Bowen, W. D., Iverson, S. J., 2003. Weaning mass affects changes in body composition and food intake in harbour seal pups during the first month of independence. *Physiol. Biochem. Zool.* 76, 418-427.
- Nielsen, J., 1995. Immunological and hematological parameters in captive harbor seals (*Phoca vitulina*). *Mar. Mamm. Sci.* 11, 314-323.
- Nunn, C. L., Gittleman, J. L., Antonvics, J., 2003. A comparative study of white blood cell counts and disease risk in carnivores. *Proc. R. Soc. Lond. B.* 270, 347-356.
- Nunn, C. L., Lindenfors, P., Pursall, E. R., Rolff, J., 2009. On sexual dimorphism in immune function. *Phil. Trans. R. Soc. B.* 364, 61-69.
- Oki, C., Atkinson, S., 2004. Diurnal patterns of cortisol and thyroid hormones in the Harbor Seal (*Phoca vitulina*) during summer and winter seasons. *Gen. Comp. Endocrinol.* 136, 289-297.
- Ortiz, R. M., Wade, C. E., Ortiz, C. L., 2000. Prolonged fasting increases the response of the Renin-Angiotensin-Aldosterone system, but not vasopressin levels, in postweaned northern elephant seal pups. *Gen. Comp. Endocrinol.* 119, 217-223.
- Ortiz, R. M., Worthy, G. A. J., 2000. Effects of capture on adrenal steroid and vasopressin concentrations in free-ranging bottlenose dolphins (*Tursiops truncatus*). *Comp. Biochem. Physiol. A.* 125, 317-324.

- Ortiz, R. M., Noren, D. P., Litz, B., Ortiz, C. L., 2001a. A new perspective on adiposity in a naturally obese mammal. *Am. J. Physiol. Endocrinol. Metab.* 281, E1347-E1351.
- Ortiz, R. M., Wade, C. E., Ortiz, C. L., 2001b. Effects of prolonged fasting on plasma cortisol and TH in postweaned northern elephant seal pups. *Am. J. Physiol. Regul. Integ. Comp. Physiol.* 280, R790-R795.
- Ortiz, R. M., Houser, D. S., Wade, C. E., Ortiz, C. L., 2003a. Hormonal changes associated with the transition between nursing and natural fasting in northern elephant seals (*Mirounga angustirostris*). *Gen. Comp. Endocrinol.* 130, 78-83.
- Ortiz, R. M., Wade, C. E., Ortiz, C. L., Talamantes, F., 2003b. Acutely elevated vasopressin increases circulating concentrations of cortisol and aldosterone in fasting northern elephant seal (*Mirounga angustirostris*) pups. *J. Exp. Biol.* 206, 2795-2802.
- Ortiz, R. M., Crocker, D. E., Houser, D. S., Webb, P. M., 2006. Angiotensin II and aldosterone increase with fasting in breeding adult male northern elephant seals (*Mirounga angustirostris*). *Physiol. Biochem. Zool.* 79, 1106-1112.
- Pitcher, K. W., 1986. Variation in blubber thickness of harbor seals in Southern Alaska. *J. Wildl. Manag.* 50, 463-466.
- Pruessner, J. C., Kirschbaum, C., Meinlschmid, G., Hellhammer, D. H., 2003. Two formulas for computation of the area under the curve represent measures of total hormone concentration versus time-dependent change. *Psychoneuroendocr.* 28, 916-931.
- Re, R.N., Kourides, I.A., Ridgway, E.C., Weintraub, B.D., Maloof, F., 1976. The effect of glucocorticoid administration on human pituitary secretion of thyrotropin and prolactin. *J. Clin. Endocrinol. Metab.* 43, 338-346.
- Riviere, J. E., Englehardt, F. R., Solomon, J., 1977. The relationship of thyroxine and cortisol to the moult of the harbor seal *Phoca vitulina*. *Gen. Comp. Endocrinol.* 31, 398-401.
- Romano, T. A., Keogh, M. J., Kelly, C., Feng, P., Berk, L., Schlundt, C. E., Carder, D. A., Finneran, J. J., 2004. Anthropogenic sound and marine mammal health: measures of the nervous and immune systems before and after intense sound exposure. *Can. J. Fish. Aquat. Sci.* 61, 1124-1134.

- Romero, L. M., Meister, C. J., Cyr, N. E., Kenagy, G. J., Wingfield, J. C., 2008. Seasonal glucocorticoid responses to capture in wild free-living mammals. *Am. J. Physiol. Regul. Integr. Comp. Physiol.* 294, R614-R622.
- Rosen, D. A. S., Renouf, D., 1997. Seasonal changes in blubber distribution in Atlantic harbor seals: indication of thermodynamic considerations. *Mar. Mamm. Sci.* 13, 229-240.
- Rosen, D. A. S., Renouf, D., 1998. Correlates of seasonal changes in metabolism in Atlantic harbour seals (*Phoca vitulina concolor*). *Can. J. Zool.* 76, 1520-1528.
- Rosen, D. A. S., Kumagai, S., 2008. Hormone changes indicate that winter is a critical period for food shortages in Steller sea lions. *J. Comp. Physiol. B.* 178, 573-583.
- Ross, P. S., Pohajdak, B., Bowen, W. D., Addison, R. F., 1993. Immune function in free-ranging harbor seal (*Phoca vitulina*) mothers and their pups during lactation. *J. Wildl. Dis.* 29, 21-29.
- Roth, J. A., Kaeberle, M. L., 1981a. Effects of *in vivo* dexamethasone administration on *in vitro* bovine polymorphonuclear leukocyte function. *Infect. Immun.* 33, 434-441.
- Roth, J. A., Kaeberle, M. L., 1981b. Evaluation of bovine polymorphonuclear leukocyte function. *Vet. Immunol. Immunopathol.* 2, 157-174.
- Saeb, M., Baghshani, H., Nazifi, S., Saeb, S., 2010. Physiological response of dromedary camels to road transportation in relation to circulating levels of cortisol, thyroid hormones and some serum biochemical parameters. *Trop. Anim. Health Prod.* 42, 56-63.
- Sangalang, G. B., Freeman, H. C., 1976. Steroids in the plasma of the gray seal, *Halichoerus grypus*. *Gen. Comp. Endocrinol.* 29, 419-422.
- St. Aubin, D. J., Geraci, J. R., 1986. Adrenocortical function in pinniped hyponatremia. *Mar. Mamm. Sci.* 2, 243-250.
- St. Aubin, D. J., Geraci, J. R., 1988. Capture and handling stress suppresses circulating levels of thyroxine (T4) and triiodothyronine (T3) in beluga whales *Delphinapterus leucas*. *Physiol. Zool.* 61, 170-175.
- Stratakis, C. A., Chrousos, G. P., 1995. Neuroendocrinology and pathophysiology of the stress system. *Ann. N. Y. Acad. Sci.* 771, 1-18.

- Swart, R. L. D., Kluten, R. M. G., Huizing, C. J., Vedder, L. J., Reijnders, P. J. H., Visser, I. K. G., Uytdehaag, F. G. C. M., Osterhaus, A. D. M. E., 1993. Mitogen and antigen induced B and T cell responses of peripheral blood mononuclear cells from the harbour seal (*Phoca vitulina*). Vet. Immunol. Immunopathol. 37, 217-230.
- Swart, R. L. D., Ross, P. S., Vedder, L. J., Timmerman, H. H., Heisterkamp, S., Loveren, H. V., Vos, J. G., Reijnders, P. J. H., Osterhaus, A. D. M. E., 1994. Impairment of immune function in harbor seals (*Phoca vitulina*) feeding on fish from polluted waters. Ambio 23, 155-159.
- Thomson, C. A., Geraci, J. R., 1986. Cortisol, aldosterone, and leucocytes in the stress response of bottlenose dolphins, *Tursiops truncatus*. Can. J. Fish. Aquat. Sci. 43, 1010-1016.
- Watson, A. D. J., Church, D. B., Emslie, D. R., Foster, S. F., 1998. Plasma cortisol responses to three corticotrophic preparations in normal dogs. Aust. Vet. J. 76, 255-257.
- Weber, K. T., 2003. A Neuroendocrine-immune interface. Herz. 28, 692-701.
- Wechsler-Reya, R. J., Monroe, J. G., 1996. Lipopolysaccharide prevents apoptosis and induces responsiveness to antigen receptor cross-linking in immature B cells. Immunol. 89, 356-362.
- Westly, H. J., Kelley, K. W., 1984. Physiologic concentrations of cortisol suppress cell-mediated immune events in the domestic pig. Proc. Soc. Exp. Biol. Med. 177, 156-164.
- Zapol, W. M., Liggins, G. C., Schneider, R. C., Qvist, J., Snider, M. T., Creasy, R. K., Hochachka, P. W., 1979. Regional blood flow during simulated diving in the conscious Weddell seal. J. Appl. Physiol. 47, 968-973.

Chapter 5:

General Conclusions

This study assessed the “health” of individual animals by quantifying circulating concentrations of hormones associated with nutritional status and fat mass (leptin), lipid and water metabolism (cortisol and aldosterone), and growth and metabolism (thyroxine and triiodothyronine) as well as circulating total and differential leukocyte counts and *in vitro* proliferation of PBMC.

5.1.1 Hormones, Body Mass, and Age

There is strong evidence supporting the importance of body mass in reproductive success and survival of pups in phocids and otariids (Baker and Fowler, 1992; Arnbom *et al.*, 1994; Kjellqvist *et al.*, 1995; Ellis *et al.*, 2000; Bowen *et al.*, 2001; Harding *et al.*, 2005). Further, changes in body condition or mass in juvenile and adult pinnipeds have been associated with changes in circulating cortisol, thyroid hormones and leptin in several species of terrestrial mammals (Concannon *et al.*, 2001; Buff *et al.*, 2002; Barboza *et al.*, 2004; Caldeira *et al.*, 2007) and pinnipeds (Ortiz *et al.*, 2003; Jeanniard du Dot *et al.*, 2009). However, in the present studies body condition index was not related to any hormones quantified in SSL pups (Chapter 3). Further, age of SSL pups during the early postnatal period, while significant was only weakly related to total and free T₄ concentrations ($r^2=0.098$). When compared to juvenile and sub-adult SSL (Rosen and Kumagai, 2008; Jeanniard du Dot *et al.*, 2009), SSL pups (Chapter 3) had two times greater total T₄ concentrations while free T₄ concentrations were half the circulating concentration found in juvenile and sub-adult SSL (Jeanniard du Dot *et al.*, 2009). Therefore, the weak relationship between age and thyroid hormone concentrations in SSL pups is likely due to the short developmental period (5-38 days) covered in the present study (Chapter 3), suggesting that a longitudinal study which covers a longer

development period is likely needed to fully explore the influence of age on circulating thyroid hormone concentrations.

The relationship of leptin concentrations and body mass in pinnipeds remains unclear. Previous studies have reported difficulty in quantifying leptin in SSL (Rosen and Kumagai, 2008) and other pinnipeds (Arnould *et al.*, 2002; Ortiz *et al.*, 2003) have been previously reported with low concentrations or no correlation to body mass or lipid stores (Ortiz *et al.*, 2001; Arnould *et al.*, 2002; Rosen and Kumagai, 2008). The present study (Chapter 3) does not clarify the role of leptin in pinnipeds. Leptin concentrations in our study were near the lower detection limit of the kit (0.5 ng/ml) similar to previous studies in phocids with similar methods (Hammond *et al.*, 2005).

BCI or body mass and nutritional status have showed seasonality in phocids (Ryg *et al.*, 1990; Beck *et al.*, 1993; Renouf *et al.*, 1993; Nilssen *et al.*, 2001; Chabot and Stenson, 2002) and otariids (Kumagai *et al.*, 2006; Mellish *et al.*, 2007; Rea *et al.*, 2007; Williams *et al.*, 2007; Jeanniard du Dot *et al.*, 2008). Juvenile HS did not show seasonal variation in cortisol or aldosterone, nor did body mass or composition affect the response to an acute stressor in the present study (Chapter 4). Sexually mature HS have been reported to display seasonal variation in body mass, lipid stores (Rosen and Renouf, 1997; Mellish *et al.*, 2007), energy intake (Rosen and Renouf, 1998), and resting metabolic rate (Rosen and Renouf, 1998). However, seasonal variation was not found in body mass, energy intake or blubber stores in previous studies of juvenile HS (Pitcher, 1986; Markussen and Ryg, 1990). Therefore, the lack of seasonality may be due to HS in the present study being immature (Chapter 4). Given these findings exogenous ACTH administration should be applied to sexually mature HS to determine if seasonality is present in the response to an acute stressor which was not present in the juvenile HS (Chapter 4).

Overall, the findings of this study indicate that mass and body condition did not influence endocrine or immune profiles of juvenile HS and SSL pups. However, caution should be taken given the short developmental period (SSL pups) and low variation in body condition and mass (HS and SSL pups) covered in the present study. When the

findings of the present study are compared to published studies on SSL and HS, there does appear to be an effect of age, at least as classified as mature vs. immature animals in thyroid hormones and total and differential leukocyte counts.

5.1.2 Total and Differential Leukocyte Counts

Across mammalian species the number of circulating total white blood cell counts has been shown to increase with body mass in adults (Nunn *et al.*, 2003; Nunn *et al.*, 2009). However, total and differential WBC counts were not related to body mass or composition in juvenile HS (Chapter 4). Compared to wild adult HS (Ross *et al.*, 1993; Morgan *et al.*, 1998) baseline WBC, neutrophil, and eosinophil counts were lower while lymphocyte and monocytes counts were higher in the present study (Chapter 4). Differences in total and differential white blood cell counts between immature and mature mammals have been reported across taxa including carnivore (Nunn *et al.*, 2009). Therefore, differences in total and differential leukocyte counts observed in the present study (Chapter 4) may be related to age of the HS. Nielsen (1995) found that total WBC count decreased in association of an increase in neutrophils (%) with a concurrent decrease in lymphocytes (%) with age (0.75 - 17.5 years of age) of HS. Therefore, the decrease in WBC was driven primarily by a reduced number of circulating lymphocytes. When HS pups were monitored during the early postnatal period, pups showed an initial increase in WBC count, which was maintained until at least weaning (Ross *et al.*, 1993; Ross *et al.*, 1994). The influence of age on total and differential leukocytes in HS differs from the findings for SSL pups in the present study (Chapter 2). In SSL pups, total WBC counts showed a decrease between 5 and 38 days of age. Unlike HS, the decrease in total WBC count was primarily driven by a decrease in neutrophils while no change was observed in lymphocyte counts. These findings along with previous studies on older pups (6 to 10 months of age; Mellish *et al.*, 2006) and juveniles (1 to 3 years; Mellish *et al.*, 2006) indicate that young SSL pups have a higher WBC count at birth followed by a postnatal decrease remaining below the levels observed in juveniles. Previous studies on SSL have not reported differential leukocyte counts and therefore further comparisons in

changes of neutrophil and lymphocyte populations are not possible. However, it is likely that the reported change in total WBC counts in SSL is driven predominately by changes in circulating neutrophil counts. These findings are suggestive that SSL and HS pups show different patterns of changes in the circulating populations of leukocytes during the early postnatal period. Previous studies on HS pups and the current study on SSL pups (Chapter 2) have explored postnatal changes in PBMC function by *in vitro* mitogen stimulation. However, to date no studies on either HS or SSL have explored changes to the innate immune response. Given the changes associated with age in circulating numbers of neutrophils observed in both HS and SSL, future studies should explore immunological functions of neutrophils, such as phagocytic and respiratory burst activity and the effect of age.

Studies focusing on neutrophils may provide informative not only in regards to functional changes associated with age or maturity but also in regards to the effects of an acute stressor on neutrophils. Elevated cortisol concentrations have been associated with capture and handling methods routinely used during blood collection in wild mammals (Engelhard *et al.*, 2002; Romero *et al.*, 2008) including in the present study (Chapter 2, Chapter 3). GC have been shown to negatively affect some components of the immune system (Marketon and Glaser, 2008) with administration of ACTH or cortisol producing characteristic changes in circulating populations of leukocytes (Roth and Kaeberle, 1981a; Roth and Kaeberle, 1981b). During an acute stress response circulating neutrophil counts increase rapidly with a concurrent decrease in lymphocyte and monocytes (Dhabhar *et al.*, 1995; Dhabhar *et al.*, 1996; Cattet *et al.*, 2003). Similarly an increase in total WBC, neutrophil and neutrophil to lymphocyte ratio was associated with an increase in cortisol concentration following ACTH administration in juvenile HS (Chapter 4). These findings demonstrate that juvenile HS response to an acute stressor is typically mammalian exhibiting a stress leukogram. However, the number of circulating lymphocytes and the *in vitro* proliferation of PBMC did not change in association with rookery disturbance and associated handling in SSL pups (Chapter 2) or following administration of ACTH in juvenile HS (Chapter 4). The innate immune response of

neutrophils was not explored and given the increase in circulating neutrophils associated with an acute stressor in juvenile HS (Chapter 4), future studies should include assays to assess the function of neutrophils including phagocytic and respiratory burst activity.

5.1.3 PBMC Proliferation

All animals in the present studies responded to *in vitro* stimulation with ConA and LPS 055:B5 indicating peripheral T and B cells are capable of responding to an antigenic challenge. The use of two mitogens that preferentially stimulate either T cells (Barta and Barta, 1993) or B cells (Wechsler-Reya and Monroe, 1996) allowed the assessment of both T and B cell populations. *In vitro* proliferation of B cells was lower than T cells in both SSL (Chapter 2) and HS (Chapter 4) similar to previous observations in HS (Levin *et al.*, 2005). The lower proliferation of B cells is likely the result of a lower number of B cells than T cells in the peripheral blood as observed in other mammals (Byrne *et al.*, 2000; Faldyna *et al.*, 2001). A decrease in T cell proliferation but no change in B cell proliferation was associated with increasing age during the postnatal period in SSL (Chapter 2). The observed age related changes in cell-mediated immunity were not associated with a decrease in circulating lymphocyte or monocytes during the postnatal period in SSL (Chapter 2). The observed changes in T cell proliferation associated with age in SSL (Chapter 2) and lower proliferation of B cells compared to T cells in SSL pups (Chapter 2) and juvenile HS (Chapter 4) may be related to the ratio of T and B cells in circulation; however, I did not quantify the number of circulating T and B cells. The ratio of T and B cells has not been explored in pinnipeds and will need to be determined in future studies. These studies will depend on the cross reactivity of commercially available antibodies or development of species-specific antibodies.

In addition to establishing normal ranges for T and B cells in pinnipeds, the application of immunophenotyping lymphocyte subpopulations would give further insight into leukocyte trafficking during an acute stress response. GC have been shown to negatively affect the immune system (Marketon and Glaser, 2008) with administration of ACTH or cortisol producing characteristic changes in circulating populations of

leukocytes often referred to the stress leukogram. As part of the stress leukogram, lymphocyte counts decrease in mammals (Dhabhar *et al.*, 1995; Dhabhar *et al.*, 1996). Juvenile HS displayed the classic stress leukogram 240 min after administration of exogenous ACTH (Chapter 4), while no change in lymphocyte counts was associated with the rookery disturbance and handling of SSL pups (Chapter 2). In rats the reduction in circulating lymphocytes associated with restraint stress was due to a decrease in both circulating B and T cells; however, the reduction of B cells was greater (Dhabhar *et al.*, 1995). Future studies to determine the number and proportion of circulating subpopulations of lymphocytes would provide insight into the relationship between changes observed in the number of circulating lymphocytes and the differences or changes in proliferation of T and B cells in response to age and acute stressors.

5.1.4 Effect of Handling and Sampling

In the present study, cortisol and all forms of thyroid hormones quantified decreased when regressed against elapsed time since arrival on rookery in wild SSL pups (Chapter 3). The decrease in total and free T₄ and total T₃ is likely the result of the initial rookery disturbance and associated increased activity of pups. However, the elapsed time is by necessity later in the day and therefore the decrease in thyroid hormones may not solely be related to the rookery disturbance but rather the result of increasing temperatures as the day progressed. Further, cortisol concentrations influenced total and differential leukocytes counts in juvenile harbor seal (HS) (Chapter 4). These findings highlight the importance of assessing the impact of potential stressors such as rookery disturbance and animal handling when sampling pinnipeds.

5.2 Literature Cited

- Arnbom, T., Fedak, M. A., Rothery, P., 1994. Offspring sex ratio in relation to female size in southern elephant seals, *Mirounga leonina*. Behav. Ecol. Sociobiol. 35, 373-378.
- Arnould, J. P. Y., Morris, M. J., Rawlins, D. R., Boyd, I. L., 2002. Variation in plasma leptin levels in response to fasting in Antarctic fur seals (*Arctocephalus gazella*). J. Comp. Physiol. B. 172, 27-34.

- Baker, J. D., Fowler, C. W., 1992. Pup weight and survival of northern fur seals *Callorhinus ursinus*. J. Zool. Lond. 227, 231-238.
- Barboza, P. S., Hartbauer, D. W., Hauer, W. E., Blake, J. E., 2004. Polygynous mating impairs body condition and homeostasis in male reindeer (*Rangifer tarandus tarandus*). J. Comp. Physiol. B. 174, 309-317.
- Barta, O., Barta, V., 1993. Lymphocyte transformation test. In: Barta, O. (Ed.), Veterinary Clinical Immunology Laboratory. Monographs in Animal Immunology, 2. Bar-Lab, Inc Blacksburg, pp. B4-1-B4-17.
- Beck, G. G., Smith, T. G., Hammill, M. O., 1993. Evaluation of body condition in the Northwest Atlantic harp seal (*Phoca groenlandica*). Can. J. Fish. Aquat. Sci. 50, 1372-1381.
- Bowen, W. D., Ellis, S. L., Iverson, S. J., Boness, D. J., 2001. Maternal effects on offspring growth rate and weaning mass in harbour seals. Can. J. Zool. 79, 1088-1101.
- Buff, P. R., Dodds, A. C., Morrison, C. D., Whitley, N. C., McFadin, E. L., Daniel, J. A., Djiane, J., Keisler, D. H., 2002. Leptin in horses: tissue localization and relationship between periperal concentrations of leptin and body condition. J. Anim. Sci. 80, 2942-2948.
- Byrne, K. M., Kim, H. W., Chew, B. P., Reinhart, G. A., Hayek, M. G., 2000. A standardized gating technique for the generation of flow cytometry data for normal canine and normal feline blood lymphocytes. Vet. Immunol. Immunopathol. 73, 167-182.
- Caldeira, R. M., Belo, A. T., Santos, C. C., Vazques, M. I., Portugal, A. V., 2007. The effect of body condition score on blood metabolites and hormonal profiles in ewes. Sm. Rumin. Res. 68, 233-241.
- Cattet, M. R. L., Christison, K., Caulkett, N. A., Stenhouse, G. B., 2003. Physiologic responses of grizzly bears to different methods of capture. J. Wildl. Dis. 39, 649-654.
- Chabot, D., Stenson, G. B., 2002. Growth and seasonal fluctuations in size and condition of male Northwest Atlantic harp seals *Phoca groenlandica*: an analysis using sequential growth curves. Mar. Ecol. Prog. Ser. 227, 25-42.
- Concannon, P., Levac, K., Rawson, R., Tennant, B., Bensadoun, A., 2001. Seasonal changes in serum leptin, food intake, and body weight in photoentrained woodchucks. Am. J. Physiol. Regul. Integr. Comp. Physiol. 281, R951-R959.

- Dhabhar, F. S., Miller, A. H., McEwen, B. S., Spencer, R. L., 1995. Effects of stress on immune cell distribution. *J. Immunol.* 154, 5511-5527.
- Dhabhar, F. S., Miller, A. H., McEwen, B. S., Spencer, R. L., 1996. Stress-induced changes in blood leukocytes distribution: role of adrenal steroid hormones. *J. Immunol.* 157, 1638-1644.
- Ellis, S. L., Bowen, W. D., Boness, D. J., Iverson, S. J., 2000. Maternal effects on offspring mass and stage of development at birth in the harbor seal, *Phoca vitulina*. *J. Mammal.* 81, 1143-1156.
- Engelhard, G. H., Brasseur, S. M. J. M., Hall, A. J., Burton, H. R., Reijnders, P. J. H., 2002. Adrenocortical responsiveness in southern elephant seal mothers and pups during lactation and the effect of scientific handling. *J. Comp. Physiol. B.* 172, 315-328.
- Faldyna, M., Leva, L., Knotigova, P., Toman, M., 2001. Lymphocyte subsets in peripheral blood of dogs - a flow cytometric study. *Vet. Immunol. Immunopathol.* 82, 23-37.
- Hammond, J. A., Bennett, K. A., Walton, M. J., Hall, A. J., 2005. Molecular cloning and expression of leptin in gray and harbor seal blubber, bone marrow, and lung and its potential role in marine mammal respiratory physiology. *Am. J. Physiol. Regul. Integr. Comp. Physiol.* 289, R545-R553.
- Harding, K. C., Fugiwara, M., Axberg, Y., Harkonen, T., 2005. Mass-dependent energetics and survival in harbour seal pups. *Func. Ecol.* 19, 129-135.
- Jeanniard du Dot, T., Rosen, D. A. S., Trites, A. W., 2008. Steller sea lions show diet-dependent changes in body composition during nutritional stress and recover more easily from mass loss in winter than in summer. *J. Exp. Mar. Biol. Ecol.* 367, 1-10.
- Jeanniard du Dot, T., Rosen, D. A. S., Richmond, J. P., Kitaysky, A. S., Zinn, S. A., Trites, A. W., 2009. Changes in glucocorticoid, IGF-I and thyroid hormones as indicators of nutritional stress and subsequent refeeding in Steller sea lions (*Eumetopias jubatus*). *Comp. Biochem. Physiol. A.* 152, 524-534.
- Kjellqwist, S. A., Haug, T., Oritsland, T., 1995. Trends in age-composition, growth and reproductive parameters of Barents Sea harp seals, *Phoca groenlandica*. *ICES J. Mar. Sci.* 52, 197-208.

- Kumagai, S., Rosen, D. A. S., Trites, A. W., 2006. Body mass and composition responses to short-term low energy intake are seasonally dependent in Steller sea lions (*Eumetopias jubatus*). *J. Comp. Physiol. B.* 176, 589-598.
- Levin, M., De Guise, S. D., Ross, P. S., 2005. Association between lymphocyte proliferation and polychlorinated biphenyls in free-ranging harbor seal (*Phoca vitulina*) pups from British Columbia, Canada. *Environ. Toxicol. Chem.* 24, 1247-1252.
- Marketon, J. I. W., Glaser, R., 2008. Stress hormones and immune function. *Cell. Immunol.* 252, 16-26.
- Markussen, N. H., Ryg, M., 1990. Energy requirements for maintenance and growth of captive harbour seals, *Phoca vitulina*. *Can. J. Zool.* 68, 423-426.
- Mellish, J.-A. E., Calkins, D. G., Christen, D. R., Horning, M., Rea, L. D., Atkinson, S. K., 2006. Temporary captivity as a research tool: comprehensive study of wild pinnipeds under controlled conditions. *Aquat. Mamm.* 32, 58-65.
- Mellish, J.-A. E., Horning, M., York, A. E., 2007. Seasonal and spatial blubber depth changes in captive harbor seals (*Phoca vitulina*) and Steller's sea lions (*Eumetopias jubatus*). *J. Mammol.* 88, 408-414.
- Morgan, L., Kumaresan, S., Thomas, C., MacWilliams, P., 1998. Hematology and chemistry reference values for free-ranging harbor seals (*Phoca vitulina*) and the effects of hemolysis on chemistry values of captive harbor seals. *J. Zoo Wildl. Med.* 29, 394-400.
- Nielsen, J., 1995. Immunological and hematological parameters in captive harbor seals (*Phoca vitulina*). *Mar. Mamm. Sci.* 11, 314-323.
- Nilssen, K. T., Haug, T., Lindblom, C., 2001. Diet of weaned pups and seasonal variations in body condition of juvenile Barents Sea harp seals *Phoca groenlandica*. *Mar. Mamm. Sci.* 17, 926-936.
- Nunn, C. L., Gittleman, J. L., Antonvics, J., 2003. A comparative study of white blood cell counts and disease risk in carnivores. *Proc. R. Soc. Lond. B.* 270, 347-356.
- Nunn, C. L., Lindenfors, P., Pursall, E. R., Rolff, J., 2009. On sexual dimorphism in immune function. *Phil. Trans. R. Soc. B.* 364, 61-69.
- Ortiz, R. M., Noren, D. P., Litz, B., Ortiz, C. L., 2001. A new perspective on adiposity in a naturally obese mammal. *Am. J. Physiol. Endocrinol. Metab.* 281, E1347-E1351.

- Ortiz, R. M., Houser, D. S., Wade, C. E., Ortiz, C. L., 2003. Hormonal changes associated with the transition between nursing and natural fasting in northern elephant seals (*Mirounga angustirostris*). Gen. Comp. Endocrinol. 130, 78-83.
- Pitcher, K. W., 1986. Variation in blubber thickness of harbor seals in Southern Alaska. J. Wildl. Manag. 50, 463-466.
- Rea, L. D., Rosen, D. A. S., Trites, A. W., 2007. Utilization of stored energy reserves during fasting varies by age and season in Steller sea lions. Can. J. Zool. 85, 190-200.
- Renouf, D., Gales, R., Noseworthy, E., 1993. Seasonal variation in energy intake and condition of harp seals: Is there a harp seal morph? Problems for bioenergetic modelling. J. Zool. Lond. 230, 513-528.
- Romero, L. M., Meister, C. J., Cyr, N. E., Kenagy, G. J., Wingfield, J. C., 2008. Seasonal glucocorticoid responses to capture in wild free-living mammals. Am. J. Physiol. Regul., Integr. Comp. Physiol. 294, R614-R622.
- Rosen, D. A. S., Renouf, D., 1997. Seasonal changes in blubber distribution in Atlantic harbor seals: indication of thermodynamic considerations. Mar. Mamm. Sci. 13, 229-240.
- Rosen, D. A. S., Renouf, D., 1998. Correlates of seasonal changes in metabolism in Atlantic harbour seals (*Phoca vitulina concolor*). Can. J. Zool. 76, 1520-1528.
- Rosen, D. A. S., Kumagai, S., 2008. Hormone changes indicate that winter is a critical period for food shortages in Steller sea lions. J. Comp. Physiol. B. 178, 573-583.
- Ross, P. S., Pohajdak, B., Bowen, W. D., Addison, R. F., 1993. Immune function in free-ranging harbor seal (*Phoca vitulina*) mothers and their pups during lactation. J. Wildl. Dis. 29, 21-29.
- Ross, P. S., Swart, R. L. D., Visser, I. K. G., Vedder, L. J., Murk, W., Bowen, W. D., Osterhaus, A. D. M. E., 1994. Relative immunocompetence of the newborn harbour seal, *Phoca vitulina*. Vet. Immunol. Immunopathol. 42, 331-348.
- Roth, J. A., Kaeberle, M. L., 1981a. Evaluation of bovine polymorphonuclear leukocyte function. Vet. Immunol. Immunopathol. 2, 157-174.
- Roth, J. A., Kaeberle, M. L., 1981b. Effects of *in vivo* dexamethasone administration on *in vitro* bovine polymorphonuclear leukocyte function. Infect. Immun. 33, 434-441.

- Ryg, M., Smith, T. G., Oritsland, N. A., 1990. Seasonal changes in body mass and body composition of ringed seals (*Phoca hispida*) on Svalbard. *Can. J. Zool.* 68, 470-475.
- Wechsler-Reya, R. J., Monroe, J. G., 1996. Lipopolysaccharide prevents apoptosis and induces responsiveness to antigen receptor cross-linking in immature B cells. *Immunol.* 89, 356-362.
- Williams, T. M., Rutishauser, M., Long, B., Fink, T., Gafney, J., Mostman-Liwanag, H., Casper, D., 2007. Seasonal variability in Otariid energetics: Implications for the effects of predators on localized prey resources. *Physiol. Biochem. Zool.* 80, 433-443.

Appendix A. List of commercially available RIA used to quantify circulating hormone concentrations.

Hormone	Manufacturer	Catalog	Range	Sensitivity
Cortisol	Siemens	TK01	1-50 µg/dL	0.2 µg/dL
Aldosterone	Siemens	PITKAL-4	25-1200 pg/ml	11 pg/ml
Triiodothyronine (T ₃)	Siemens	TKT31	20-600 ng/dL	7 ng/dL
Thyroxin (T ₄)	Siemens	TKT41	0.5-24 µg/dL	0.25 µg/dL
Free thyroxin (T ₄)	Siemens	PITKF4-5	0.1-10 ng/dL	0.01 ng/dL
Leptin	Linco	XL-85K	1-50 ng/ml	0.5 ng/ml

Vacutainer CPT with sodium citrate (8 ml, BD 362761; 4 ml, BD)

Cell Proliferation Biotrak ELISA system, version 2 (GE Healthcare Bio-Sciences Corp., RPN 250)

Concanavalin A (Sigma Aldrich, C 04152)

Lipopolysaccharide from *Escherichia coli* 055:B5 (Sigma Aldrich, L-2880)

Appendix B. Standard Operating Protocol for Aldosterone Assays

OBJECTIVE: To determine aldosterone concentration in unextracted sera in Steller sea lions and harbor seals.

HEALTH AND SAFETY:

Proper lab attire including lab coats, gloves and safety glasses are required as needed or as indicated by MSDSs for chemicals being tested.

PERSONNEL/TRAINING/RESPONSIBILITIES:

Any lab employee or student familiar with the equipment and laboratory techniques and trained in this and any referenced SOPs may perform this procedure.

REQUIRED AND RECOMMENDED MATERIALS:

Aldosterone coat-a-count kit (Siemens cat #PITKAL-4)
VWR lab marker
(4) non coated tubes
Serum samples

PROCEDURE:

1. Allow sera to thaw and come to room temperature.
2. Label tubes.
3. Pipette 50 μ l of zero calibrator A into NSB and A tubes.
4. Pipette 50 μ l of remaining calibrator, controls.
5. Pipette serum into the appropriate tubes.
6. Add 0.250 ml 125 I Aldosterone to every tube
7. Gently mix
8. Incubate for 18 hrs at room temperature
9. Decant thoroughly, removing all visible moisture.
10. Count for 1 minute in the gamma counter.

Tubes	13. E (200 pg/ml)
1. TC Plain tube	14. "
2. "	15. F (600 pg/ml)
3. NSB Plain tube	16. "
4. "	17. G (1200 pg/ml)
5. A (0 pg/ml)	18. "
6. "	19. Control Low
7. B (25 pg/ml)	20. "
8. "	21. Control High
9. C (50 pg/ml)	22. "
10. "	23. Sample #1
11. D (100 pg/ml)	24. "
12. "	

Appendix C. Standard Operating Protocol for Cortisol Assays

OBJECTIVE: To determine cortisol concentrations in unextracted sera from Steller sea lions and harbor seals.

HEALTH AND SAFETY:

Proper lab attire including lab coats, gloves and safety glasses are required as needed or as indicated by MSDSs for chemicals being tested.

PERSONNEL/TRAINING/RESPONSIBILITIES:

Any lab employee or student familiar with the equipment and laboratory techniques and trained in this and any referenced SOPs may perform this procedure. Personnel must be trained for radioactive techniques.

REQUIRED AND RECOMMENDED MATERIALS:

Cortisol Coat-a-count kit (Siemens TK01)

Serum samples

VWR lab marker

(4) non coated tubes

PROCEDURE:

1. Allow sera to thaw and come to room temperature.
2. Label tubes.
3. Pipette 12.5 μ l of zero calibrator A into:
 - a. NSB tubes
 - b. A calibrator tube
4. Pipette 12.5 μ l of remaining calibrator and samples into the appropriate tubes.
5. Add 500 μ l of 125 I Cortisol to every tube.
6. Vortex
7. Incubate at 37°C for 45 min.
8. Decant supernatant (except TC 1, 2).
9. Count for 1 minute in a gamma counter.

Tubes	9. 5 μ g/dl (C)	18. "
1. Total Count	10. "	19. Control 5
2. "	11. 10 μ g/dl (D)	20. "
3. NSB	12. "	21. Control 6
4. "	13. 20 μ g/dl (E)	22. "
5. 0 μ g/dl (A)	14. "	23. Sample #1
6. "	15. 50 μ g/dl (F)	24. "
7. 1 μ g/dl (B)	16. "	
8. "	17. Control 4	

Appendix D. Standard Operating Protocol total thyroxine (T₄) Assays

OBJECTIVE: To determine total T₄ levels in sera from harbor seals and Steller sea lions.

HEALTH AND SAFETY:

Proper lab attire including lab coats, gloves and safety glasses are required as needed or as indicated by MSDSs for chemicals being tested.

PERSONNEL/TRAINING/RESPONSIBILITIES:

Any lab employee or student familiar with the equipment and laboratory techniques and trained in this and any referenced SOPs may perform this procedure. Personnel must be trained for RADIOACTIVE techniques.

REQUIRED AND RECOMMENDED MATERIALS:

Total T₄ RIA kit (Siemens TKT41)
 Serum samples
 VWR marker
 (4) non coated tubes

PROCEDURE:

1. Allow kit to come to room temperature.
2. Allow sera to thaw and come to room temperature.
3. Label tubes
 - a. Plain tubes- (2) Total Count; (2) NSB
 - b. Remaining tubes are coated tubes
4. Pipette 12.5 µl of zero calibrator A into:
 - a. NSB tubes
 - b. A calibrator tube
5. Pipette 12.5 µl of remaining calibrator and samples into the appropriate tubes.
6. Add 500 µl of ¹²⁵I Total T₄ to every tube.
7. Vortex
8. Incubate at 37°C for 60 min.
9. Decant supernatant (except TC 1, 2) - remaining inverted for at least 2-3 min.
10. Count for 1 minute in a gamma counter.

Tubes	9. 4 µg/dl (C)	17. Control low (B/C)
1. Total Count	10. "	18. "
2. "	11. 10 µg/dl (D)	19. Control High (E/F)
3. NSB	12. "	20. "
4. "	13. 16 µg/dl (E)	21. Sample #1
5. 0 µg/dl (A)	14. "	22. "
6. "	15. 24 µg/dl (F)	
7. 1 µ/dl (B)	16. "	
8. "		

Appendix E. Standard Operating Protocol for Free thyroxine (T₄) Assays

OBJECTIVE: To determine free T₄ levels in unextracted sera from harbor seals and Steller sea lions.

HEALTH AND SAFETY:

Proper lab attire including lab coats, gloves and safety glasses are required as needed or as indicated by MSDSs for chemicals being tested.

PERSONNEL/TRAINING/RESPONSIBILITIES:

Any lab employee or student familiar with the equipment and laboratory techniques and trained in this and any referenced SOPs may perform this procedure. Personnel must be trained for RADIOACTIVE techniques.

REQUIRED AND RECOMMENDED MATERIALS:

Total T₄ RIA kit (Siemens PITKF4-5)
Sera samples
VWR marker
(4) non coated tubes

PROCEDURE:

1. Allow kit to come to room temperature.
2. Allow sera to thaw and come to room temperature.
3. Label tubes
 - a. Plain tubes- (2) Total Count; (2) NSB
 - b. Remaining tubes are coated tubes
4. Pipette 25 µl of zero calibrator A into:
 - a. NSB tubes
 - b. A calibrator tube
5. Pipette 25 µl of remaining calibrator and samples into the appropriate tubes.
6. Add 500 µl of ¹²⁵I free T₄ to every tube.
7. Vortex
8. Incubate at 37°C for 60 min.
9. Decant supernatant (except TC 1, 2)- remaining inverted for at least 2-3 min.
10. Count for 1 minute in a gamma counter.

Tubes	9. 0.5 ng/dl (C)	18. "
1. Total Count	10. "	19. Cntrl low (B/C)
2. "	11. 1.3 ng/dl (D)	20. "
3. NSB	12. "	21. Cntrl High (F/G)
4. "	13. 2.2 ng/dl (E)	22. "
5. 0 ng/dl (A)	14. "	23. Sample #1
6. "	15. 4.8 ng/dl (F)	24. "
7. 0.1 ng/dl (B)	16. "	
8. "	17. 10 ng/dl (G)	

Appendix F. Standard Operating Protocol for Total triiodothyronine (T₃) Assays

OBJECTIVE: To determine total T₃ concentrations in unextracted sera from harbor seals and Steller sea lions.

HEALTH AND SAFETY:

Proper lab attire including lab coats, gloves and safety glasses are required as needed or as indicated by MSDSs for chemicals being tested.

PERSONNEL/TRAINING/RESPONSIBILITIES:

Any lab employee or student familiar with the equipment and laboratory techniques and trained in this and any referenced SOP may perform this procedure. Personnel must be trained for RADIOACTIVE techniques.

REQUIRED AND RECOMMENDED MATERIALS:

Total T₃ RIA kit (Siemens TKT31)
Serum samples
VWR marker

PROCEDURE:

1. Allow kit to come to room temperature.
2. Allow sera to thaw and come to room temperature.
3. Label tubes
 - a. Plain tubes- (2) Total Count; (2) NSB
 - b. Remaining tubes are coated tubes
4. Pipette 50 ul of zero calibrator A into:
 - a. NSB tubes
 - b. A calibrator tube
5. Pipette 50 ul of remaining calibrator and samples into the appropriate tubes.
6. Add 500 ul of ¹²⁵I total T₃ to every tube.
7. Vortex
8. Incubate at 37°C for 2 hours.
9. Decant supernatant (except TC 1,2)- remaining inverted for at least 2-3 min.
10. Count for 1 minute in a gamma counter.

Tubes	9. 20 ng/dl (B)	17. 600 ng/dl (F)
	10. "	18. "
1. Total Count	11. 50 ng/dl (C)	19. Control low (B/C)
2. "	12. "	20. "
3. NSB	13. 100 ng/dl (D)	21. Control high (E/F)
4. "	14. "	22. "
5. 0 ng/dl (A)	15. 200 ng/dl (E)	23. Sample #1
6. "	16. "	24. "

Appendix G. Standard Operating Protocol for Leptin Assays

OBJECTIVE: To determine Leptin levels in sera from harbor seals.

HEALTH AND SAFETY:

Proper lab attire including lab coats, gloves and safety glasses are required as needed or as indicated by MSDSs for chemicals being tested.

PERSONNEL/TRAINING/RESPONSIBILITIES:

Any lab employee or student familiar with the equipment and laboratory techniques and trained in this and any referenced SOPs may perform this procedure.

REQUIRED AND RECOMMENDED MATERIALS:

- Leptin RIA kit (Linco XL-85K)
- Serum samples (1:1)
- VWR marker
- Glass tubes

PROCEDURE:

Day 1

1. Allow kit to come to room temperature.
2. Allow sera to thaw and come to room temperature.
3. Label tubes.
4. Pipette 150 μ l of assay buffer:
 - a. NSB tubes (tube 3,4)
 - b. Bo (tube 5,6)
5. Pipette 50 μ l:
 - a. Standards (tubes
 - b. Quality controls (tubes
 - c. Samples
6. Pipette 50 μ l Multispecies Leptin antibody to tubes 5 on.
7. Vortex, cover, and incubate overnight (20-24 hrs) at 4°C water bath

Day 3

8. Pipette 50 μ l of 125 I-Human Leptin to all tubes.
9. Vortex, cover, and incubate overnight (20-24 hrs) at 4°C water bath.

Day 4

10. Add 500 μ l cold (4°C) Precipitating Reagent to all tubes except total count (tubes 1,2)
11. Vortex and incubate 20 minutes at 4°C.
12. Centrifuge at 4°C, 3000 g, 30 min
13. Decant supernatant (except TC 1,2)
14. Count all tubes on gamma counter for 1 minute

Tubes

1. Total Count
2. “
3. NSB
4. “
5. 0 ng/ml (A)
6. “
7. 1 ng/ml (B)
8. “
9. 2 ng/ml (B)
10. “
11. 5 ng/ml (D)
12. “
13. 10 ng/ml (E)
14. “
15. 20 ng/ml (F)
16. “
17. 50 ng/ml (G)
18. “
19. Control Low
20. “
21. Control High
22. “
23. E110
24. “

Appendix H. Standard Operating Protocol for cell preparation tubes (CPT™)

OBJECTIVE: To isolate and freeze peripheral blood mononuclear cells (PBMC) for lymphocyte proliferation assay.

HEALTH AND SAFETY:

Proper lab attire including lab coats, gloves and safety glasses are required as needed or as indicated by MSDSs for chemicals being tested.

PERSONNEL/TRAINING/RESPONSIBILITIES:

Any lab employee or student familiar with the equipment and laboratory techniques and trained in this and any referenced SOPs may perform this procedure.

REQUIRED AND RECOMMENDED MATERIALS:

2 (8ml) Vacutainer CPT with sodium citrate (BD 362761) per animal
 PBS w/out Ca^{2+} / Mg^{2+} (Celgro, 21-040-CV)
 Freezing Media (RPMI-20 + 10% DMSO)
 Cryovials
 Sterile transfer Pipettes
 Gloves
 70% ETOH
 VWR Marker

PROCEDURE:

1. Thaw freezing media. Keep cold.
2. Collect blood into the CPT tube using standard techniques (invert tube several times to mix blood and anticoagulant).
3. After collection, tube can be kept upright at room temperature for up to two hours, before centrifugation.
4. Remix blood sample immediately prior to centrifugation (gently invert 8-10x)
5. Centrifuge tube for 20 min, 20°C, 1500 x g. Once centrifuged tube can be stored upright at 4°C for up to 24 hrs until processed further.
6. From this point on should be done under the sterile hood.
7. Using the sterile pipette carefully remove the top clear layer of liquid (only 1-2 ml) and discard.
8. Transfer the remaining contents i.e. the “fuzzy” layer above the gel and remaining clear solution into a 15 ml centrifuge tube containing 8mls of PBS.
9. Mix cells and PBS by gentle inversion-5x.
10. Centrifuge for 8 min at 950 rpm, 20°C.
11. Decant the supernatant without disturbing cell pellet.
12. Add 1 ml freezing media and mix thoroughly by repeated pipetting.
13. Transfer cells and freezing media into cryovials.
14. Place cryovials in freezing chamber and store at -80°C until the next day.
15. Remove cryovials from freezing chamber and place in liquid N₂ dewar.

Appendix I. Standard Operating Protocol for Sodium Heparin Tubes

OBJECTIVE: To freeze WBC for future lymphocyte isolation and proliferation assay.

10 ml Na- Heparin green top tube (GT)

Freezing Media

Cryovials

Transfer Pipettes

Gloves

70% ETOH

VWR Marker

PROCEDURE:

1. Thaw freezing media. Keep cold.
2. Collect blood into the GT tube using standard techniques (invert tubes 4-5x to mix blood and anticoagulant).
3. After collection, tube can be upright and chilled until centrifugation.
4. Remix blood sample immediately prior to centrifugation (gently invert 4-5x).
5. Centrifuge tube for 10 min at 950 rpm.
6. From this point on should be done under the sterile hood.
7. Using the individually wrapped sterile pipette carefully remove the top clear layer of plasma without disturbing the white buffy layer.
8. Transfer plasma into cryovials and store at -80°C.
9. Gently transfer the buffy coat (i.e. "fuzzy" layer above RBC) into 1.2 ml cryovials.
10. Using a new individually wrapped sterile pipette add 1 ml freezing media to the WBC cryovial and mix thoroughly by gently inverting the cryovial containing the WBC and Freezing media.
11. Place cryovials in freezing chamber and store at -80°C until the next day.
12. Remove cryovials from freezing chamber and place in liquid N₂ dewar.

Appendix J: Standard Operating Protocol for PBMC Proliferation Assay

I. Thawing Cells

1. Remove vial from liquid N₂ and thaw quickly in 37°C water bath.
2. Transfer contents of vial into 10 mls of RPMI-10 Media.
3. Centrifuge for 10 min, 1000 rpm, 20°C (Wash 1).
4. Aspirate supernatant without disturbing cell pellet.
5. Re-suspend cells; add 10 ml RPMI-10.
6. Allow cells to sit at room temperature for 2 hours.
7. Centrifuge for 10 min, 1000 rpm, 20°C (Wash 2).
8. Re-suspend cells; add 5 ml RPMI-10
9. Transfer 50 µl of cells to eppendorf tube with 200 µl RPMI-10,
10. Count cells.
11. Bring volume up to 10 ml Centrifuge for 10 min, 1000 rpm, 20°C (Wash 3).
12. Re-suspend cells in C-RPMI-10 at a final concentration of 4×10^6 cells/ml;
(2×10^5 cells/50 µl)

II. Preparing Mitogens

Reagents Needed:

C- RPMI-10

Stock mitogen(s) 1 mg/ml

Concanavalin A (Sigma, cat # C 04152)

LPS 055:B5 (Sigma, cat # L-2880)

13. Based on the experiment, determine which concentration of mitogen requires the highest volume.
14. Using this volume as the minimum, add an additional amount.
15. Continue with this process until the lowest concentration needed is made.

III. Plating Cells

Place the following in the hood:

Eight channel pipette

Pipette tips

Sterile boat (one for each animal)

16. Label the plate (s).
17. With the eight channel pipette, aliquot the mitogens (50 µl/well) into their respective wells according to the schematic.
18. Once all of the mitogens have been plated, precede to aliquot the cells into their respective wells using 50 µl/well.
19. Cover plates when finished and observe cells under the microscope.
20. Place in 37°C, 5% CO₂ incubator for 48 hrs.

Make sure enough CO₂ is left in the tanks and that the metal tray at the bottom of the incubator contains enough sterile water.

IV. Labeling the Cells with BrdU

Allow all reagents to reach room temperature before assay.

Mix samples and all reagents prior to use.

Avoid excessive foaming of reagents.

21. Dilute the BrdU labeling reagent (bottle 1) 1:100 with C-RPMI-10 [100 μ M].
22. Add 10 μ l/well BrdU labeling solution.
23. Re-incubate cell cultures for an additional 18 hrs at 37°C, 5% CO₂.
24. Centrifuge the plate at 1500 rpm for 10 min, 20°C.
25. Remove media by tapping.
26. Dry cells in drying oven (60°C) for 1 hr.
27. Add 200 μ l/well fixative and incubate for 30 min at room temperature.
28. Remove fixative by tapping.
29. Add 200 μ l/well blocking buffer.
30. Remove blocking buffer by tapping.
31. Add 100 μ l/well peroxidase-labelled anti-BrdU working solution
(1:100 antibody dilution solution)
32. Incubate for 90 min at room temperature.
33. Remove antibody conjugate by tapping.
34. Rinse wells three times with 200 μ l/well washing solution.
35. Immediately dispense 100 μ l of room temperature equilibrated TMB substrate into all wells.
36. Cover the plate and mix at room temperature until color development is sufficient (5-30 min).
37. Add 25 μ l of 1M sulphuric acid (H₂SO₄) to each well in order to stop reaction.
38. Determine the optical density in the plate reader at 450 nm within 5 min.

Cell Count:

1. Label 0.5 mL sterile eppendorf tubes for each animal.
2. Using a sterile 1ml pipette add 200 μ L C-RPMI-5 into each eppendorf tube.
3. Using a sterile eppendorf tip add 50 μ L cells into the appropriate tube.
4. Mix well.
5. The counts are done non-sterilely so remove just the eppendorf tubes from the hood.
6. In a 96 well flat-bottom plate mix 50 μ L of the cells/media with 50 μ L Trypan blue.
7. Gives a total dilution of 1:10.
8. Using a hemacytometer and microscope count four grid squares for each animal.

Calculations:

Total cells/4= (Ave cell) (10) (10⁴) = # cells/ml

Appendix K. Solutions

1M sulphuric acid (H_2SO_4)

560 μl concentrated sulphuric acid to 8 ml of **ice-cold** distilled water.
Carefully mix and make up to 10 ml with water

All steps for the following solutions should be performed using sterile techniques in a sterile hood. Wash hood with 70% alcohol before and after use.

C-RPMI-10 Media

RPMI 1640 w/out L-glutamine (Gibco, 21870-092)
50 ml FBS (Hyclone, SH3007103HI)
5 ml Glutamate-Penicillin-Streptomycin (Gibco, 10378-016)
5 ml 100 mM Sodium Pyruvate (Gibco, 11360-070)
5 ml 100x Nonessential Amino Acids (Gibco, 11140-050)
5 ml 1M HEPES (Gibco, 15630-080)
500 μl 0.05 M 2-mercaptoethanol (Sigma, M7522-100ML)
Store at 4°C

0.05 M 2-mercaptoethanol

20 μl 14.3 M 2-mercaptoethanol (Sigma, # M-7522)
5.6 ml RPMI-10

RPMI-10 Media

RPMI 1640 w/out L-glutamine (Gibco, 21870-092)
50 ml FBS (Hyclone, SH3007103HI)
5 ml Glutamate-Penicillin-Streptomycin (Gibco, 10378-016)
Store at 4°C

LPS 055:B5 stock [1mg/ml]

For every mg of LPS 055:B55 add 1 ml of C-RPMI-10. Mix thoroughly and aliquot into sterile 1 μl eppendorf and store at -80°C .

ConA stock [1mg/ml]

For every mg of Con A add 1 ml of C-RPMI-10. Mix thoroughly and aliquot into sterile 500 μl eppendorf and store at -80°C .

Freezing Media

RPMI-10 Media
20% FBS (Hyclone, SH3007103HI)
10% DMSO (Sigma)
Mix thoroughly and aliquot 5-10 ml aliquots into sterile 15 ml conical tubes and store at -80°C .